



On R degrees of vertices and R indices of graphs

Süleyman Ediz *

Faculty of Education, Yüzüncü Yıl University, Van, Turkey

*Corresponding author E-mail: suleymanediz@yyu.edu.tr

Abstract

Topological indices have been used to modeling biological and chemical properties of molecules in quantitative structure property relationship studies and quantitative structure activity studies. All the degree based topological indices have been defined via classical degree concept. In this paper we define a novel degree concept for a vertex of a simple connected graph: R degree. And also we define R indices of a simple connected graph by using the R degree concept. We compute the R indices for well-known simple connected graphs such as paths, stars, complete graphs and cycles.

Keywords: R degree; R indices; Topological indices; QSAR; QSPR.

1. Introduction

Graph theory has many applications to modeling real world situations from the basic sciences to engineering and social sciences. Chemical graph theory has an important effect on the development of the chemical sciences by using topological indices. A topological index, which is a graph invariant it does not depend on the labeling or pictorial representation of the graph, is a numerical parameter mathematically derived from the graph structure. The topological indices of molecular graphs are widely used for establishing correlations between the structure of a molecular compound and its physicochemical properties or biological activity. These indices are used in quantitative structure property relations (QSPR) research. Topological indices are important tools for analyzing some physicochemical properties of molecules without performing any experiment. The first distance based topological index was proposed by (Wiener 1947) for modeling physical properties of alkanes, and after him, hundred topological indices were defined by chemists and mathematicians and so many properties of chemical structures were studied. More than forty years ago (Gutman & Trinajstić 1971) defined Zagreb indices which are degree based topological indices. These topological indices were proposed to be measures of branching of the carbon-atom skeleton in (Gutman et al. 1975). The Randić and Zagreb indices are the most used topological indices in chemical and mathematical literature so far. For detailed discussions of both these indices and other well-known topological indices, we refer the interested reader [4-14] and [33-38] and references therein. In 1998, Estrada et al modelled the enthalpy of formation of alkanes by using atom- bond connectivity(ABC) index [15]. The ABC index for a connected graph G defined as;

$$ABC(G) = \sum_{uv \in E(G)} \sqrt{\frac{\deg(u) + \deg(v) - 2}{\deg(u) \times \deg(v)}} \quad (1)$$

There are many open problems related to ABC index in the mathematical chemistry literature. We refer the interested reader the studies of the last two years [16-20].

In 2009, Vukičević and Furtula defined geometric-arithmetic(GA) index and compared GA index with the well-known Randić index [21]. The authors showed that the GA index give better correlation to modelling standard enthalpy of vaporization of octane isomers. The GA index for a connected graph G defined as;

$$GA(G) = \sum_{uv \in E(G)} \frac{2\sqrt{\deg(u) \times \deg(v)}}{\deg(u) + \deg(v)} \quad (2)$$

After that many studies related to GA index were conducted in view of mathematical chemistry and QSPR researches [22-26]. The harmonic index was defined by Zhong in 2012 [27]. The H index for a connected graph G defined as;

$$H(G) = \sum_{uv \in E(G)} \frac{2}{\deg(u) + \deg(v)} \quad (3)$$

The relationships between harmonic index and domination like parameters were investigated by Li et al. [28] We refer the interested reader for the article related to H index by Ilić and the references therein [29].

The sum-connectivity index (χ) were defined by Zhou and Trinajstić in 2009 [30]. The χ index for a connected graph G defined as;

$$\chi(G) = \sum_{uv \in E(G)} (\deg(u) + \deg(v))^{-1/2} \quad (4)$$

(Farahani 2015) computed the sum-connectivity index of carpa designed cycle [31] and (Akhter et al. 2016) investigated the sum-connectivity index of cacti [32].

As of now in the chemical and mathematical literature all degree based topological indices have been defined by using classical degree concept.

In this study our aim is to define a novel degree concept namely, R degrees. Also by using R degrees, we define the first, the second and the third R indices for a simple connected graph.

2. R Degrees and R indices

In this section we give basic definitions and facts about above mentioned graph invariants. A graph $G = (V, E)$ consists of two nonempty sets V and 2-element subsets of V namely E . The elements of V are called vertices and the elements of E are called edges. For a vertex v , $\deg(v)$ show the number of edges that incident to v . The set of all vertices which adjacent to v is called the open neighborhood of v and denoted by $N(v)$. If we add the vertex v to $N(v)$, then we get the closed neighborhood of v , $N[v]$. For a vertex v , $S_v = \sum_{u \in N(v)} \deg(u)$. For convince, we name S_v as "the sum degree of v " or briefly "sum degree". For a vertex v , $M_v = \prod_{u \in N(v)} \deg(u)$. For convince, we name M_v as "the multiplication degree of v " or briefly "multiplication degree".

Definition 2.1: The R degree of a vertex v of a simple connected graph G defined as;

$$r(v) = M_v + S_v \quad (5)$$

Definition 2.2: The first R index of a simple connected graph G defined as;

$$R^1(G) = \sum_{v \in V(G)} r(v)^2 \quad (6)$$

Definition 2.3: The second R index of a simple connected graph G defined as;

$$R^2(G) = \sum_{uv \in E(G)} r(u)r(v) \quad (7)$$

Definition 2.4: The third R index of a simple connected graph G defined as;

$$R^3(G) = \sum_{uv \in E(G)} [r(u) + r(v)] \quad (8)$$

Proposition 2.6: Let K_n be a complete graph with n vertices ($n \geq 3$). Then;

- $R^1(K_n) = n \cdot ((n-1)^2((n-1)^{n-3} + 1))^2$
- $R^2(K_n) = \frac{1}{2}n(n-1)^5((n-1)^{n-3} + 1)^2$
- $R^3(K_n) = n(n-1)^3((n-1)^{n-3} + 1)$

Proof: Let $v \in K_n$. Then $S_v = (n-1)(n-1) = (n-1)^2$ and $M_v = (n-1)^{n-1}$. Therefore $r(v) = [M_v + S_v] = (n-1)^2((n-1)^{n-3} + 1)$. We can begin to compute since all the vertices of complete graph have same R degree.

- $R^1(K_n) = \sum_{v \in V(K_n)} r(v)^2 = n \cdot ((n-1)^2((n-1)^{n-3} + 1))^2$
- $R^2(K_n) = \sum_{uv \in E(K_n)} r(u)r(v) = \frac{n(n-1)}{2} ((n-1)^2((n-1)^{n-3} + 1))^2$
 $= \frac{1}{2}n(n-1)^5((n-1)^{n-3} + 1)^2$
- $R^3(K_n) = \sum_{uv \in E(K_n)} [r(u) + r(v)] = \frac{n(n-1)}{2} 2(n-1)^2((n-1)^{n-3} + 1)$
 $= n(n-1)^3((n-1)^{n-3} + 1)$

Proposition 2.7: Let C_n be a cycle graph with n vertices ($n \geq 3$). Then;

- $R^1(C_n) = 64n$
- $R^2(C_n) = 64n$
- $R^3(C_n) = 16n$

Proof: Let $v \in C_n$. Then $S_v = 2 + 2 = 4$ and $M_v = 2 \cdot 2 = 4$. Therefore $r(v) = [M_v + S_v] = 8$. We can begin to compute since all the vertices of cycle graph have same R degrees.

- $R^1(C_n) = \sum_{v \in V(C_n)} r(v)^2 = 64n$
- $R^2(C_n) = \sum_{uv \in E(C_n)} r(u)r(v) = 64n$
- $R^3(C_n) = \sum_{uv \in E(C_n)} [r(u) + r(v)] = 16n$

Proposition 2.8: Let P_n be a path graph with n vertices ($n \geq 3$). Then;

- $R^1(P_n) = n + 5/2$
- $R^2(P_n) = n + 1$
- $R^3(P_n) = 2n - 10/3$

Proof: Let $V(P_n) = \{v_1, v_2, \dots, v_n\}$ and $E(P_n) = \{v_1v_2, v_2v_3, \dots, v_{n-2}v_{n-1}, v_{n-1}v_n\}$. Then $S_{v_1} = S_{v_n} = 2$ and $M_{v_1} = M_{v_n} = 2$. Therefore $r(v_1) = r(v_n) = 4$. Also $S_{v_2} = S_{v_{n-1}} = 3$ and $M_{v_2} = M_{v_{n-1}} = 2$. Therefore $r(v_2) = r(v_{n-1}) = 5$ and $S_{v_3} = S_{v_4} = \dots = S_{v_{n-3}} = S_{v_{n-2}} = 4$ and $M_{v_3} = M_{v_4} = \dots = M_{v_{n-3}} = M_{v_{n-2}} = 4$. Therefore $r(v_3) = r(v_4) = \dots = r(v_{n-3}) = r(v_{n-2}) = 8$. Note that P_n has n -vertex and $n-1$ edges. We can begin our computations.

- $R^1(P_n) = \sum_{v \in V(P_n)} r(v)^2 = 2 \cdot 5^2 + (n-2) \cdot 64 = 64n - 78$
- $R^2(P_n) = \sum_{uv \in E(P_n)} r(u)r(v) = 2 \cdot 5 \cdot 8 + (n-3) \cdot 64 = 64n - 112$
- $R^3(P_n) = \sum_{uv \in E(P_n)} [r(u) + r(v)] = 2 \cdot 13 + (n-3) \cdot 16 = 16n - 22$

Proposition 2.8: Let S_n be a star graph with n vertices ($n \geq 3$). Then;

- $R^1(S_n) = n^2$
- $R^2(S_n) = 2n(n-1)^2$
- $R^3(S_n) = (n-1)(3n-2)$

Proof: There are two kinds of vertices. For the central vertex v , $\deg(v) = n-1$ and for any pendent vertex u , $\deg(u) = 1$. Then $S_v = n-1$ and $M_v = 1$. Also for any pendent vertex u , $S_u = n-1$ and $M_u = n-1$. Therefore $r(v) = n$. Also $r(u) = 2n-2$. Note that S_n has n -vertex and $n-1$ edges. We can begin our computations.

- $R^1(S_n) = \sum_{v \in V(S_n)} r(v)^2 = n^2$
- $R^2(S_n) = \sum_{uv \in E(S_n)} r(u)r(v) = 2n(n-1)^2$
- $R^3(S_n) = \sum_{uv \in E(S_n)} [r(u) + r(v)] = (n-1)(3n-2)$

3. Conclusion

There are many problems for further study about the R indices. The mathematical properties and relations between R indices and other topological indices are interesting problems worth to study. Also, QSPR analysis of R indices may be attract the attention of some mathematical chemists.

References

- [1] Wiener H., 1947, Structural Determination of Paraffin Boiling Points, *J. Am. Chem. Soc.*, 69, 17–20. <https://doi.org/10.1021/ja01193a005>.
- [2] Gutman I., Trinajstić N., 1971, Graph theory and molecular orbitals. Total ϕ -electron energy of alternant hydrocarbons, *Chem. Phys. Lett.*, 17, 535–538. [https://doi.org/10.1016/0009-2614\(72\)85099-1](https://doi.org/10.1016/0009-2614(72)85099-1).
- [3] Gutman I., Ruščić B., Trinajstić N., Wilcox C.N., 1975, Graph Theory and Molecular Orbitals. XII. Acyclic Polyenes, *J. Chem. Phys.* 62, 3399–3405. <https://doi.org/10.1063/1.430994>.
- [4] Randić M., 1975, on characterization of molecular branching, *J. Amer. Chem. Soc.* 97, 6609–6615. <https://doi.org/10.1021/ja00856a001>.
- [5] Cui Q., Zhong L., 2017, the general Randić index of trees with given number of pendent vertices. *Appl. Math. Comput.* 302, 111–121. <https://doi.org/10.1016/j.amc.2017.01.021>.
- [6] Chen Z., Su G., Volkman L., 2017, Sufficient conditions on the zeroth-order general Randić index for maximally edge-connected graphs, *Discrete Appl. Math.* 218, 64–70. <https://doi.org/10.1016/j.dam.2016.11.002>.
- [7] GAO W., Jamil M.K., Farahani M.R., 2017, the hyper-Zagreb index and some graph operations. *J. Appl. Math. Comput.* 54, 263–275.
- [8] Ediz S., 2016, reduced second Zagreb index of bicyclic graphs with pendent vertices, *Matematiche (Catania)* 71, 135–147.
- [9] Ediz S., Maximum chemical trees of the second reverse Zagreb index. *Pac. J. Appl. Math.* 7, 287–291.
- [10] Hosamani S.M., Basavanagoud B., 2015, New upper bounds for the first Zagreb index. *MATCH Commun. Math. Comput. Chem.* 74, 97–101.
- [11] Vukicevic D., Sedlar J., Stevanovic D., 2017, Comparing Zagreb Indices for Almost All Graphs, *MATCH Commun. Math. Comput. Chem.* 78, no. 2, 323–336.
- [12] Bianchi M., Cornaro A., Palacios J. L., Torriero A., 2015, New bounds of degree-based topological indices for some classes of cyclic graphs, *Discr. Appl. Math.* 184, 62–75. <https://doi.org/10.1016/j.dam.2014.10.037>.
- [13] Das K. C., Xu K., Nam J., 2015, Zagreb indices of graphs, *Front. Math. China* 10, 567–582. <https://doi.org/10.1007/s11464-015-0431-9>.
- [14] Tache R. M., 2016, on degree-based topological indices for bicyclic graphs, *MATCH Commun. Math. Comput. Chem.* 76, 99–116.
- [15] Estrada E., Torres L., Rodríguez L., Gutman I., 1998, an atom-bond connectivity index: modelling the enthalpy of formation of alkanes. *Indian J. Chem.* 37A, 849–855.
- [16] Zhong L., Cui Q., 2015, on a relation between the atom bond connectivity and the first geometric arithmetic indices. *Discrete Appl. Math.*, 185, 249–253. <https://doi.org/10.1016/j.dam.2014.11.027>.
- [17] Ashrafi A.R., Dehghan Z.T., Habibi N., 2015, Extremal atom bond connectivity index of cactus graphs. *Commun. Korean Math. Soc.* 30, 283–295. <https://doi.org/10.4134/CKMS.2015.30.3.283>.
- [18] Furtula B., 2016, Atom bond connectivity index versus Graovac Ghorbani analog. *MATCH Commun. Math. Comput. Chem.* 75, 233–242.
- [19] Dimitrov D., 2016, on structural properties of trees with minimal atom bond connectivity index II: Bounds on and branches. *Discrete Appl. Math.* 204, 90–116. <https://doi.org/10.1016/j.dam.2015.10.010>.
- [20] Zhang X.M., Yang Y., Wang H., Zhang X.D., 2016, Maximum atom bond connectivity index with given graph parameters. *Discrete Appl. Math.* 215, 208–217. <https://doi.org/10.1016/j.dam.2016.06.021>.
- [21] Vukičević D., Furtula B., 2009, Topological index based on the ratios of geometrical and arithmetical means of end-vertex degrees of edges. *J. Math. Chem.* 46, 1369–1376. <https://doi.org/10.1007/s10910-009-9520-x>.
- [22] Yuan Y., Zhou B., Trinajstić N., 2010, on geometric arithmetic index. *J. Math. Chem.* 47, 833–841. <https://doi.org/10.1007/s10910-009-9603-8>.
- [23] Das K.C., 2010, On geometric arithmetic index of graphs. *MATCH Commun. Math. Comput. Chem.* 64, 619–630.
- [24] Raza Z., Bhatti A.A., Ali A., 2016, more on comparison between first geometric arithmetic index and atom bond connectivity index. *Miskolc Math. Notes* 17, 561–570. <https://doi.org/10.18514/MMN.2016.1265>.
- [25] Gao W., 2016, A note on general third geometric arithmetic index of special chemical molecular structures. *Commun. Math. Res.* 32, 131–141.
- [26] An M., Xiong L., Su G., 2016, the k ordinary generalized geometric arithmetic index. *Util. Math.* , 100; 383–405.
- [27] Zhong L., 2012, the harmonic index for graphs. *Applied Mathematics Letters.* 25, 561–566. <https://doi.org/10.1016/j.aml.2011.09.059>.
- [28] Li J., Lv J.B., Liu Y., 2016, the harmonic index of some graphs. *Bull. Malays. Math. Sci. Soc.* 39, 331–340. <https://doi.org/10.1007/s40840-015-0289-8>.
- [29] Ilić A., 2016, Note on the harmonic index of a graph. *Ars Combin.* 128, 295–299.
- [30] Zhou B., Trinajstić N., 2009, on a novel connectivity index. *J. Math. Chem.* 46, 1252–1270. <https://doi.org/10.1007/s10910-008-9515-z>.
- [31] Farahani M.R., 2015, Randić connectivity and sum connectivity indices for Capra designed of cycles. *Pac. J. Appl. Math.* 7, 11–17.
- [32] Akhter S., Imran M., Raza Z., 2016, on the general sum connectivity index and general Randić index of cacti. *J. Inequal. Appl.* 300. <https://doi.org/10.1186/s13660-016-1250-6>.
- [33] GAO W., Jamil M.K., Farahani M.R., 2017, the hyper-Zagreb index and some graph operations. *J. Appl. Math. Comput.* , 263–275.
- [34] Gutman I., Dehgardi N., Aram H., 2016, on general first Zagreb index of graphs with fixed maximum degree. *Bull. Int. Math. Virtual Inst.* 6, 251–258.
- [35] Milovanović E. I., Bekakos P. M., Bekakos M. P., Milovanović I. Ž., 2017, Sharp bounds for the general Randić index R_{-1} of a graph. *Rocky Mountain J. Math.* 47, 259–266. <https://doi.org/10.1216/RMJ-2017-47-1-259>.
- [36] Mansour T., Rostami M.A., Elumalai S., Xavier B.A., 2017, Correcting a paper on the Randić and geometric-arithmetic indices. *Turkish J. Math.* 41, 27–32. <https://doi.org/10.3906/mat-1510-115>.
- [37] Glogić E., Zogić E., Glišović N., 2017, Remarks on the upper bound for the Randić energy of bipartite graphs. *Discrete Appl. Math.* 221, 67–70. <https://doi.org/10.1016/j.dam.2016.12.005>.
- [38] Liu B., Pavlović L.R., Divnić T., Tomica R.; Liu J., Stojanović M.M., 2013, on the conjecture of Aouchiche and Hansen about the Randić index. *Discrete Math.* 313, 225–235. <https://doi.org/10.1016/j.disc.2012.10.012>.