

# Alive failures behind the windows of cancer therapy

Muhammad Torequul Islam \*

Department of Pharmacy, Southern University Bangladesh, Mehedibag (Chittagong)-4000, Bangladesh. & Postgraduate Program in Pharmaceutical Sciences, Federal University of Piauí, Teresina (Piauí)-64.049-550, Brazil

\*Corresponding author E-mail: [rbiotufpi.br@gmail.com](mailto:rbiotufpi.br@gmail.com)

## Abstract

The multiscope process, cancer is attributable from various genes. Eventually, cancer is a complicated disease with unconstrained intercalation and impacts on the physiological system. Therefore, an ideal cancer therapy must be like a multi-edged sword. Broadly, currently, available cancer therapies are the cytoprotective, inhibitors of oncogenes, correctors, and cell destructors. Doubtless, cancer therapists are most frequently handling apoptosis and autophagy inducers, targeting of tumor suppressor genes, epigenetic and immune therapies. However, each therapy has a number of challenges yet to be resolved. This revision is aimed to find out some important points, depicting till the date, how successful we are and what are the failures behind those modes of therapeutic strategies.

**Keywords:** Apoptosis; Autophagy; Cancer Therapy; Challenges; Tumor Suppressor Gene.

## 1. Introduction

The most fearful disease, cancer is still ruling crudely throughout the world. The variations in cell types and adequate sources of occurrence are the key points of continuing challenges of cancer. The selection of therapy in cancers mainly depends on- origin (i.e. - source and tissue types) and stages, as well as patients pathophysiology. Up to date, a number of anticancer agents have been introduced with single and/or multiple activity pathways. Some of them can be used in combination and are followed by a step-wise treatment (Islam 2016a).

Surgery, chemo- and radiotherapy are more commonly attainable by the populations in comparison to other cancer treatment strategies. Each strategy has unique postulation on its mechanism(s). For an example, most of the chemotherapeutic agents and radiotherapy act through chronic induction of reactive oxygen species (ROS) on cancerous cells (Islam 2016a). In recent years, some genetic materials synthesis inhibitors, anti-angiogenesis agents, apoptosis and autophagy inducers, immunotherapy, targeting epigenetic alteration correction and tumor suppressor genes have gained much attention. However, a controlled and effective action of each treatment and an overall managing power of cancer stages are great challenges along with the avoidance of a secondary cancer induction chance. Unfortunately, the success of each treatment strategy, after a moment has been raised questions for safety, efficacy and efficiency, resistance as well as applicability in other cancer types.

Cell death is essential for life, due to it plays critical roles in regulating embryonic development, maintaining tissue homeostasis, controlling immune function, tumor suppression and infection resistance. Although, cell death eliminates unfit cells, but there must be a balance between cell death and cell proliferation (Baig et al. 2016). Notably, cancer therapy constitutes growth inhibition and/or cell death. Thus, the reduction of cell mass is a reduction of physiological processes. It is doubtless that, a reduction in immune cells has been always dangerous. Moreover, every cancer

treatment imparts a shunting effect on the immune system (Islam 2016a).

This revision focuses on an improved understanding of the negative sides (may be considered as challenges) of some commonly targeting anticancer pathways, these include apoptosis, autophagy, tumor suppressor genes, epigenetics and immunotherapy.

## 2. Autophagy

Macroautophagy (also known as autophagy) is a conserved eukaryotic composed of cells catabolic pathway that causes degradation of cellular organelles and other macromolecules via lysosomal activity as part of a recycling and protective process to maintain composed of cells fitness in a basal state as well as during stress (Kimmelman 2011). Although, autophagy is important to maintain composed of cells homeostasis, disruption of autophagy can lead to disease, including neurodegeneration, atherosclerosis and cancer (Choi et al. 2013). Generally, the functions of autophagy are dynamic with both tumor-suppressive and pro-tumorigenic roles, which depend on multiple factors, including tumor stage, cellular context, and site of origin (tissue). It is evident that, tumor-suppressive selective autophagy pathways can mitigate oncogenic signals, and conversely selective autophagic pathways may support tumor maintenance and progression (Galluzzi et al. 2015).

Moreover, autophagy loss may cause tumor initiation, where it supports the transformation to invasive cancers (Qu et al. 2003). In this case, loss of function of the autophagy machinery plays an important role. On the other hand, inhibition of autophagy leads to an accumulation of ROS, increases DNA damage, and mitochondrial defects, thus an implication of tumorigenesis (White 2015). In a recent study, in the mouse model, it has been reported that p62 accumulation upon autophagy loss can contribute to tumorigenesis (Inami et al. 2011). It may be due to an over-expression of p62 promotes oxidative stress and tumor growth (Mathew et al. 2009).

Nowadays, it is clear that tumor suppression via cellular senescence is another important mechanism in autophagy, a program of

permanent arrest of the cell division cycle that can be induced by cells in response to oncogenes in order to prevent malignant transformation (Pérez-Mancera et al. 2014). GATA4, a master regulator of the cellular senescence program has been found to turn over by p62-mediated selective autophagy (Coppé et al. 2010). Thus, a transient inhibition of autophagy may lead to an accumulation of GATA4 in the cellular senescence program. It should be noted that, cellular senescence in autophagy, if needed or not is yet to be resolved (Gewirtz 2013).

In some studies, autophagy has been found to link with an elevation of RAS-driven cancers initiation and growth (Kim et al. 2011; Lock et al. 2011). Moreover, autophagy inhibition has been evident to exert an anti-tumor effect in multiple cancer types through both cells autonomous and non-autonomous mechanisms (Thorburn 2014; Galluzzi et al. 2015; White 2015).

Although, from a mechanistic standpoint, the role of autophagy in supporting tumor proliferation is complex, but it may be due to an increase in metabolism and biosynthetic need in rapidly dividing cells in a tumor microenvironment (Kimmelman 2015). Genomic instability leads a high degree of protein misfolding (Kirkin et al. 2009). Therefore, selective autophagy is needed for the cancer cells, especially those who are suffering from adequate angiogenesis and correct protein folding events.

However, role of selective autophagy in promoting or maintaining cancer cell survival may be due to the maintaining capacity of appropriate levels of signaling complexes or degrading pro-apoptotic proteins. In a study, it has been reported that the selective autophagy pathway is important for regulating active Src levels in tumor cells to promote survival following the loss of focal adhesion kinase (FAK) signaling (Sandilands et al. 2011), which is evident for signaling to promote cell adhesion, invasion, proliferation, and survival (Thomas and Brugge 1997). Generally, Src proteins are frequently over-expressed and activated in solid cancers (Irby et al. 1999), while FAK is a critical binding partner of Src at focal adhesions where its activity is regulated and directed (Sulzmaier et al. 2014). However, FAK deletion lead to a transition of active Src from an oncogenic driver to an overactive kinase, a toxin and the cancer cells were adapted a selective autophagic pathway for degradation of overactive Src (Sandilands et al. 2012). Moreover, dysregulation of ferritinophagy may lead to some diseases (Sun et al. 2015; Zhang 2015).

### 3. Apoptosis

Apoptotic cell death is widely considered as a positive process that both prevent and treat cancer. However, it can also cause unwanted effects and even promote cancer. In some literatures, it has been suggested that too much or too little apoptosis may implicate various diseases, including neurodegeneration aeration and autoimmunity (Mattson 2000; Nagata 2010). On the other hand, inhibition of apoptosis is evident to promote cancer and blunting therapeutic responses (Letai 2008).

Apoptotic cells can actively promote the proliferation of surrounding cells and as a physiological event, this may enable apoptotic cells within a tissue to control their replacement by normal turnover or as a healing response. The ultimate result is the extensive tissue damage through the activating mitogen signaling pathway (Huh et al. 2004; Perez-Garijo et al. 2004; Ryoo et al. 2004). Apoptotic cells can stimulate the proliferation of stem cells in a Caspase-dependent manner (Li et al. 2010).

Prostaglandin E2 (PGE2), a key mediator in apoptosis-induced proliferation in mammalian systems. During apoptosis, caspases cleave and activate calcium-independent phospholipase A2 (iPLA2, also known as PLA2G6) and increase in production of arachidonic acid, which is converted to PGE2 via cyclooxygenase 1 (COX1) and COX2 (also known as PTGS1 and PTGS2) (Atsumi et al. 1998). Thus, apoptotic tumour cells have an important role in tumour regrowth and repopulation after certain therapy such as radiotherapy. This may be due to caspase-3- and p53-dependent manner, probably through the production of PGE2

(Huang et al. 2011). Moreover, production of PGE2 by apoptotic tumour cells is reported to promote chemoresistance by stimulating cancer stem cell proliferation (Kurtova et al. 2015). Generally, PGE2 has pleiotropic functions such as it may promote proliferation as well as skew immune responses towards a tumour promoting, anti-inflammatory phenotype (Zelenay et al. 2015). Thus, via PGE2 pathway apoptotic tumour cells can exert an immunosuppressive effect.

Proliferating cells constantly compete with one another for nutrients. The losers may suffer contribute an apoptotic cell death. The p53, a key tumour suppressor protein might contribute to this effect as (Kruiswijk et al. 2015). Thus, the cells deficient with p53 proteins may proliferate and facilitate the accumulation of genetic lesions that lead to cancer (Bondar and Medzhitov 2010; Marusyk et al. 2010). Moreover, apoptosis is also evident to kill healthy cells (Jeffers et al. 2003; Villunger et al. 2003; Garrison et al. 2008; Michalak et al. 2009).

It has been thought that about one million cells in our bodies undergo apoptosis every second. To be mentioned that, the apoptotic cells are efficiently engulfed and destroyed by phagocytic cells. In this regard, Caspase-dependent events help to recruit the phagocytes at the apoptosis point by releasing 'find-me' signals, and promote the engulfment of them by exposing 'eat-me' signals (Arandjelovic and Ravichandran 2015). Find-me signals such as lipid lysophosphatidylcholine (LPC), nucleotides such as ATP, the proteins fractalkine (FKN; furthermore, known as CX3-C motif chemokine ligand 1 (CX3CL1)) and lactotransferrin (LTF) may exert oncogenic functions through pleiotropic effects. FKN can stimulate angiogenesis and hypoxia-induced proliferation of prostate cancer cells and can enhance oncogenic ERBB2 receptor signaling (Tardaguila and Manes 2013; Tang et al. 2015). Interestingly, adenosine (a degradation product of extracellular ATP), can be oncogenic, supporting tumour growth, angiogenesis and immune escape (Spychala 2000).

Apoptosis can promote tumorigenesis through the recruitment and activation of phagocytic macrophages at the tumour site (Gregory and Pound 2011). Additionally, tumour-associated macrophages (TAMs) may come along with them (Noy and Pollard 2014). It is also evident that, apoptotic cells can promote tumorigenesis in a non-cell-autonomous manner (Ford et al. 2015). Besides promoting tumour growth, apoptotic cells may also facilitate metastatic tumour progression. Furthermore, through massive cell death, leading to extensive efferocytosis (clearance of dead cells), promoted TAM infiltration, stimulation of a wound-healing cytokine response and increase in metastasis, implicate the clearance of apoptotic cells in cancer progression (Stanford et al. 2014).

Controlling of caspase activity is another important fact, as the level of caspase activity required to kill a cell is not so high (Rehm et al. 2006). Therefore, the Caspase-dependent apoptotic cell death is under critically controlled environment. In a study, sub-lethal doses of tumor necrosis factors (TNF)-related apoptosis-inducing ligand (TRAIL) (and FAS/CD95/APO1 ligand) were found to lead Caspase-dependent mutations and genomic instability in surviving cells (Lovric and Hawkins 2010). Moreover, Caspase-activated DNase (CAD, also known as DFFB), under an apoptosis failure condition caused a limited CAD activity, thus the DNA damage (Enari et al. 1998).

Generally, stresses at sub-lethal doses can engage mitochondrial outer membrane permeability (MOMP) in limited numbers of mitochondria without killing the cell called minority MOMP, triggers a sub-lethal caspase activity (similar to TRAIL) and causes DNA damage and genome instability in a CAD-dependent manner. A repeated engagement of sub-lethal stress was proven to promote transformation and tumorigenesis in this way (Ichim et al. 2015). However, mitotic arrest has been found to link in this pathway (Colin et al. 2015). On the other hand, transgenic expression of pro-apoptotic BAX has been found to promote lymphomagenesis - characterized by genomic instability by suppressing the co-expression of BCL-2 (Luke et al. 2003). Moreover, not only CAD, but also caspase-3-dependent release of endonuclease G (ENDOG) from the mitochondria was also found to promote

radiation-induced DNA damage and transformation (Liu et al. 2015).

The mixed lineage leukemia gene MLL (also known as KMT2A) encodes a histone-methylating enzyme that functions as an epigenetic regulator. The MLL locus is highly susceptible to breakage and rearrangement, which can generate oncogenic MLL fusion proteins, lacking methyltransferase activity (Rao and Dou 2015). Rearrangements in MLL are recurrent oncogenic drivers in various leukaemias, including acute myelogenous leukemia (AML), myelodysplastic syndromes (MDS) and acute lymphoblastic leukemia (ALL) (Gole and Wiesmuller 2015). Otherwise, failed apoptosis causes caspase- and CAD-dependent break points in the MLL gene, thereby promoting Oncogenic rearrangements are also evident in surviving cells (Betti et al. 2000; Hars et al. 2006). The latter situation may induce DNA damage under inflammatory conditions and promotes acquired resistance to apoptosis-inducing anticancer therapies (Trinchieri 2012).

Till date, various non-apoptotic roles have been ascribed to almost all proteins classically viewed as apoptotic (Hyman and Yuan 2012; Hardwick and Soane 2013; Kilbride and Prehn 2013). Some of them function as oncogenic. For an example, BCL-2 is thought to regulate calcium homeostasis, metastasis and autophagy (Bonneau et al. 2013; Pedro et al. 2015; Choi et al. 2016).

On the other hand, many human tumour types are selectively sensitive to TRAIL-induced apoptosis (Dimberg et al. 2013) and they can gain some advantage from expressing TRAIL receptors (TRAILRs). TRAILR signal can promote cancer independently of its role in canonical apoptosis signaling (Von Karstedt et al. 2015). TRAIL receptor 2 (TRAILR2) signal can promote invasion, proliferation and migration, independent of its apoptotic function but dependent on PI3K signaling. In some studies, on failed apoptosis, TRAILR signaling was reported to lead Caspase-dependent cleavage of RHO-associated protein kinase 1 (ROCK1), activating RHO GTPase and causing membrane blebbing and cell migration (Somasekharan et al. 2013).

Moreover, various non-apoptotic, pro-oncogenic functions were described in FAS signaling, including stimulation of proliferation and migration (Peter et al. 2015). However, not only pro-proliferative effects, FAS signaling can also exert pro-survival functions. Inflammatory process coming from various sources, including apoptosis, associated with non-regulated, necrotic cell death can have both tumours promoting and tumour inhibitory effects (Vakkila and Lotze 2004; Grivennikov et al. 2010).

#### 4. Tumor suppressor genes

In 1969, Knudson first predicted the existence of tumor suppressor genes (TSGs) (Friend et al. 1986). In cancer, the inactivation of one copy of a TSG will generally need to be followed by the loss of the remaining copy of the gene. A precancerous cell would only enjoy an advantage once it loses both functional copies of a TSG that had been suppressing growth (Seshadri et al. 1987). TSGs, heterozygous loss of function can be associated with reduced gene dosage and tumorigenesis via haploinsufficiency (Cook and McCaw 2000; Quon and Berns 2001). TSGs can be silenced, such as epigenetic mechanisms, or changes in mutation frequency, such as those that occur in hyper mutator phenotypes, as TSGs undergo alteration more frequently than the oncogenes. In this sense, it is harder for a drug to target TSGs, other than oncogenes.

Till date a well-described TSGs include genes in pathways are Wnt/APC (adenomatous polyposis coli gene [APC], AXIN1, and CDH1); apoptosis/cell cycle (cyclin-dependent kinase inhibitor 2A [CDKN2A], tumor p53 [TP53], RB1, TRAF7, and CASP8); chromatin modification (ARID1A/B/2, ASXL1, ATRX, CREBBP, KDM5C, KDM6A, MEN1, MLL2/3, SETD2, ten-eleven translocation-2 [TET2], WT1, and BAP1); DNA damage repair (ataxia telangiectasia mutated [ATM], ataxia telangiectasia and Rad3 related [ATR], BRCA1/2, mutL homolog 1 [MLH1], and MSH2/6); hedgehog (PTCH1); Notch (FBXW7 and NOTCH1); phosphoinositide 3-kinase (PI3K)/AKT/mammalian target of ra-

pamycin (mTOR) (PIK3R1, phosphatase and tensin homolog [PTEN], and TSC1); Ras (CEBPA, von Hippel-Lindau [VHL], and NF1); transforming growth factor- $\beta$  (SMAD2/4); and transcriptional regulation (GATA3 and RUNX1).

Among them, mutations in p53 (encoded by TP53 gene), are the most frequent genetic alterations in cancer (Freed-Pastor and Prives 2012), as it accounts in 30% to 50% of human cancers (Cerami et al. 2012). The p53 is evident to exert a dominant-negative effect when mutated. It should be noted that, this protein (p53) is not a cell-surface protein or an enzyme, thus the difficulty to target with antibodies or enzyme inhibitors (Kenzelmann Broz and Attardi 2010; Wang et al. 2011).

The anticancer drug, tenovin-1 and tenovin-6 are evident to affect p53 posttranslational modification by inhibiting the protein deacetylation activities of sirtuins, thereby stabilization of p53 (Lain et al. 2008). However, the nuclear export inhibitors such as leptomycin B, an inhibitor of the nuclear export protein CRM1, are able to increase local p53 protein levels (Mutka et al. 2009), while the nutlin, another type of anticancer drug is used to target protein-protein interactions between p53 and MDM2 (Michaelis et al. 2011). All these cases can accumulate p53 and lead to drug resistance.

Another major fact is the efficacy of such as type of treatment. For an example, treatments with viral vectors are not able to achieve the necessary efficiency of transduction by p53 within tumors to be curative (Schuler et al. 1998; Swisher et al. 1999). Moreover, repeat administration may be hampered by host immune reactions to the virus vectors. A modified approach, tumor-specific replication-competent oncolytic viruses are evident to bind and inactivate p53. Although, the viruses can only replicate within (and kill) cells lacking functional p53, but low efficiency of delivery and non-specificity in expression limit their application (Nemunaitis et al. 2000; Lamfers et al. 2002).

Notably, mutations in p53 can up-regulate the expression of platelet-derived growth factor receptor- $\beta$  (PDGFR $\beta$ ), which in turn capable to cause invasion and metastasis (Weissmueller et al. 2014). On the other hand, vaccines containing multiple p53 peptides are able to generate a T-helper type, I response, although the responses have not yet been potent enough to be clinically beneficial (Leffers et al. 2009). In another approach used dendritic cells loaded with human leukocyte antigen class, I p53 peptides, reports to induce changes in immune regulatory mechanisms along with a strong immune suppressive effect (Schuler et al. 2014).

The second messenger PIP3 activates target proteins, such as the kinases phosphoinositide-dependent kinase-1 (PDK1) and AKT1/2/3. AKT then phosphorylates as many as 20 pro-growth targets relevant to cancer, including those activating the cell cycle, preventing apoptosis, and promoting cell growth via the kinase mTOR (Puc et al. 2005; Manning and Cantley 2007). Hyperactivation of this (PI3K/AKT/mTOR) pathway resulting from inactivation of PTEN is, at least in part, similar to the sequelae of oncogenic alterations elsewhere in the pathway, such as epidermal growth factor receptor amplification or mutation, human epidermal growth factor receptor-2 (HER2) amplification, PIK3CA (the gene encoding the catalytic subunit of PI3K) mutation, or AKT1/2 mutation (Keniry and Parsons 2008). Targeting PTEN has a complexity of feedback networks. For an example, inhibition of mTOR with agents such as rapamycin is effective in attenuating signaling, but it relieves feedback inhibition of other upstream components such as insulin, insulin-like growth factor receptor, human epidermal growth factor receptor (HER)-3, and HER-4, which can then signal through other branches of the pathway such as fork head box O (FOXO)-dependent transcription. Therefore, a combined inhibition of AKT, together with agents inhibiting HER kinases or with inhibitors of receptor tyrosine kinase stabilization by heat shock protein 90, is necessary to truly shut down signaling, as AKT inhibition alone is not able to achieve (Chandarlapaty et al. 2011; Tao et al. 2014).

In cancer cells with an impaired DNA damage repair pathway, the cell becomes addicted to another DNA damage repair pathway. The gene poly (ADP-ribose) polymerase 1 (PARP1) is involved in

non-homologous end joining (NHEJ), homologous recombination (HR), and base excision repair (BER). However, PARP1 inhibition is theoretically specific for BRCA-mutated cells (Vodenicharov et al. 2000; Bryant et al. 2005), thus limiting the application of this kind of targeted therapy. In a study, mutations in PTEN have been reported to sensitize cells to PARP1 inhibition. It has been thought that, this may be due to a down-regulation of RAD51, a critical HR gene (Gupta et al. 2009).

Radiotherapy, alone or followed by some other therapies such as surgery, is evident to cause double-strand DNA breaks (Shall et al. 2011). Moreover, PARP inhibitors appear to have activity in combination with radiotherapy (Lee et al. 2007) and also chronic lymphocytic leukemias with ATM mutations that are sensitive to cytotoxic agents (Weston et al. 2010), this may also create a chance of secondary cancer.

In some instances, specificity is crucial, such as PARP inhibitors as well as several other synthetic lethal strategies targeting DNA repair proteins are evident to cause inhibition of DNA polymerases (Martin Set al. 2010). To be noted that, when TSGs undergo homozygous deletion, the region of deletion can be quite broad, and usually covers several neighboring genes. These passenger deletions may cause nearby gene deletion, including housekeeping genes, essential for cell survival (Muller et al. 2012).

## 5. Epigenetic therapy

The genetic pathways in cancer are straightforward, while reversibility and numerous unclear talks are the plugged-in epigenetic pathways. In a recent revision, Islam (2016b) has pointed out a number of facts, including the challenges of epigenetic cancer therapy, found behind the spotlight. A summary of the revision has been plugged in here. Epigenetic events are believed to occur early in cancer development (first hit for tumorigenesis). In a study, it has been suggested that, an incomplete epigenetic resetting in an environment changing frequently can adaptively co-evolve with plasticity or maternal effects, thus the trans-generational epigenetic inheritance spanning is a possibility in biological systems. Otherwise, environmentally-occupied altered trans-generational epigenetic reprogramming has been signified in inherited diseases. Therefore, in the infrequent environmental changes relative to the generation are rescued for the incomplete epigenetic resetting. In the recent years, it has been demonstrated that the genetic and epigenetic mechanisms are not separate events in cancer. They intertwine and take advantage of each other. Being a short-term and reversible event, epigenetic change may be considered as a sub- or primary stage of genetic events. Therefore, identification of the 'first cancerous hit' and correct to the point will be a perfect epigenetic treatment strategy. It is because; the epigenetic change may turn to the normal, non-cancerous stage.

On the other hand, epigenetic dysregulation is reported to change in the pattern of gene expression, activating the tumor promoting, while silencing the TSGs. Till date, several agents have been approved by the food and drug administration (FDA) for the treatment of hematologic and malignancies, including DNA methyltransferases (DNMT) and histone deacetylase (HDAC) inhibitors, despite of their limited success.

Myelosuppression, including neutropenia and thrombocytopenia, as well as nausea and vomiting is the reported toxicities in nucleoside DNMT inhibitors. Moreover, these are cytotoxic thus the chance of a second cancer is higher in the treatment of DNMT. The nucleoside analogues cause inhibition of synthesis of DNA by forming covalent complexes with the DNMT, cause their delectation, thus the reversal of methylation patterns. The DNMT, 5-azacitidine and decitabine have low efficacy and non-specificity in their activity, while S110 is more stable and may allow prolong drug exposure time, and CP-4200 has potent cytotoxic activity.

On the other hand, the HDAC inhibitors activate both intrinsic and extrinsic apoptotic pathways and regulate the activity of TSGs (e.g. - p53 and p73). These kinds of HDAC inhibitors have proteasomal degradation, ROS generation and mitochondrial outer

membrane potential losing capacity along with the inhibition of cell differentiation and growth. These are mainly anti-inflammatory drugs, and the levels and targeted activity are quite complicated.

HDAC alone cannot express the hyper-acetylated gene. Moreover, combination therapy consisting of DNMT and HDAC used in hematological cancers has been found less efficacy for solid tumors. Otherwise, the tumor microenvironment (TME) in multiple aspects of cancer progression, particularly therapeutic resistance decreases drug penetration, confers proliferatives and antiapoptotic advantages to surviving cells, facilitates resistance without causing genetic mutations and epigenetic changes.

Histone acetyltransferases (HATs) that install acetyl groups onto lysine residues of cellular proteins such as histones, transcription factors, nuclear receptors, and enzymes have been shown to play a role in a number of diseases, including cancers. Till date, several HAT inhibitors, like bi-substrate inhibitors, natural product derivatives, small molecules, and protein-protein interaction inhibitors, have been developed, despite some undesired properties like anti-oxidant activity, reactivity, instability, low potency, or lack of selectivity between HAT subtypes and other enzymes. HATs have various cellular substrates ranging from histones and transcription factors for enzymes and nuclear receptors. The catalytic mechanisms of HAT activity in relation to enzyme kinetics of small-molecule HAT inhibitors are still poorly understood.

In conclusion, the crucial challenges yet to be resolved in epigenetic cancer therapy are pointed under: clinical, laboratory development, cell biology, chemistry, target selection and toxicology.

## 6. Immunotherapy

Although, immunotherapy in cancer by these days experienced remarkable advances (Rajasagi et al. 2014), but a number of crucial challenges are yet to be resolved such as – effects of pre-installation; efficiency and specificity; onset and duration of actions; host responsiveness (as successive treatments reduce immune power; patient's pathophysiological conditions, therefore, perception); impacts on secondary cancers; effects on resistant cancer cells in a particular or combination therapy and so on. In fact, cancer is believed to originate via multiple and complex pathways (Islam 2016a).

## 7. Conclusion

The dreadful effects in cancer may be due to: massive reduction of the number of cells (including immune cells) by killing effects of cancer treatments; mal and mass production of toxins by the rapidly proliferating abnormal cells; abnormal physiological function in place of its (cancerous cell) own duty; effects on neighboring normal cells (i.e. - hypoxia, ROS, nutrients, and other toxic metabolites); reduction of the body immune power; and overall physiological outrageous networking.

Although, progress in recent years in cancer research is remarkable, but the translation of basic cancer research findings into successful therapies is a long journey. A steady progress in an effective treatment strategy is an aspiration of the cancer therapists. Understanding each fact clearly, following to a sturdy research on its pharmacology can bring novel strategies/compounds with anti-cancer potential to the clinic. Therefore, more research is needed to avoid controversial or unclear talks, prior to proceed on with a particular cancer therapeutic strategy.

## 8. Conflict of interest

There is no conflict of interest at any point of view.

## References

- [1] Arandjelovic S, Ravichandran KS. 2015. Phagocytosis of apoptotic cells in homeostasis. *Nat. Immunol.* 16:907-917. <https://doi.org/10.1038/ni.3253>.
- [2] Atsumi G, Tajima M, Hadano A, Nakatani Y, Murakami M, Kudo I. 1998. Fas-induced arachidonic acid release is mediated by Ca<sup>2+</sup>-independent phospholipase A2 but not cytosolic phospholipase A2, which undergoes proteolytic inactivation. *J. Biol. Chem.* 273:13870-13877. <https://doi.org/10.1074/jbc.273.22.13870>.
- [3] Baig S, Seevasant I, Mohamad J, Mukheem A, Huri HZ, Kamarul T. 2016. Potential of apoptotic pathway-targeted cancer therapeutic research: Where do we stand? *Cell Death Dis.* 7:e2058. <https://doi.org/10.1038/cddis.2015.275>.
- [4] Betti CJ, Villalobos MJ, Diaz MO, Vaughan AT. 2003. Apoptotic stimuli initiate MLL–AF9 translocations that are transcribed in cells capable of division. *Cancer Res.* 63:1377-1381.
- [5] Bondar T, Medzhitov R. 2010. P53-mediated hematopoietic stem and progenitor cell competition. *Cell Stem Cell* 6:309-322. <https://doi.org/10.1016/j.stem.2010.03.002>.
- [6] Bonneau B, Prudent J, Popgeorgiev N, Gillet G. 2013. Non-apoptotic roles of Bcl-2 family: the calcium connection. *Biochim. Biophys. Acta* 1833:1755-1765. <https://doi.org/10.1016/j.bbamcr.2013.01.021>.
- [7] Bryant HE, Schultz N, Thomas HD, Parker KM, Flower D, Lopez E, et al. 2005. Specific killing of BRCA2-deficient tumours with inhibitors of poly (ADP-ribose) polymerase. *Nature* 434:913-917. <https://doi.org/10.1038/nature03443>.
- [8] Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, et al. 2012. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2:401-404. <https://doi.org/10.1158/2159-8290.CD-12-0095>.
- [9] Chandralapaty S, Sawai A, Scaltriti M, Rodrik-Outmezguine V, Grbovic-Huezo O, Serra V, et al. 2011. AKT inhibition relieves feedback suppression of receptor tyrosine kinase expression and activity. *Cancer Cell* 19:58-71. <https://doi.org/10.1016/j.ccr.2010.10.031>.
- [10] Choi AMK, Ryter SW, Levine B. 2013. Autophagy in human health and disease. *N. Engl. J. Med.* 368:651-662. <https://doi.org/10.1056/NEJMra1205406>.
- [11] Choi S, Chen Z, Tang LH, Fang Y, Shin SJ, Panarelli NC, et al. 2016. Bcl-xL promotes metastasis independent of its anti-apoptotic activity. *Nat. Comm.* 7:10384. <https://doi.org/10.1038/ncomms10384>.
- [12] Colin DJ, Hain KO, Allan LA, Clarke PR. 2015. Cellular responses to a prolonged delay in mitosis are determined by a DNA damage response controlled by Bcl-2 family proteins. *Open Biol.* 5:140156. <https://doi.org/10.1098/rsob.140156>.
- [13] Cook WD, McCaw BJ. 2000. Accommodating haploinsufficient tumor suppressor genes in Knudson's model. *Oncogene* 19:3434-3438. <https://doi.org/10.1038/sj.onc.1203653>.
- [14] Coppé J-P, Desprez P-Y, Krtolica A, Campisi J. 2010. The senescence-associated secretory phenotype: the dark side of tumor suppression. *Ann. Rev. Pathol.* 5:99-118. <https://doi.org/10.1146/annurev-pathol-121808-102144>.
- [15] Dimberg LY, Anderson CK, Camidge R, Behbakht K, Thorburn A, Ford HL. 2013. On the TRAIL to successful cancer therapy? Predicting and counteracting resistance against TRAIL-based therapeutics. *Oncogene* 32:1341-1350. <https://doi.org/10.1038/ncr.2012.164>.
- [16] Enari M, Sakahira H, Yokoyama H, Okawa K, Iwamatsu A, Nagata S. 1998. A caspase-activated DNase that degrades DNA during apoptosis, and its inhibitor ICAD. *Nature* 391:43-50. <https://doi.org/10.1038/34112>.
- [17] Ford CA, Petrova S, Pound JD, Voss JJ, Melville L, Paterson M, et al. 2015. Oncogenic properties of apoptotic tumor cells in aggressive B cell lymphoma. *Curr. Biol.* 25:577-588. <https://doi.org/10.1016/j.cub.2014.12.059>.
- [18] Freed-Pastor WA, Prives C. 2012. Mutant p53: one name, many proteins. *Genes Dev.* 26:1268-1286. <https://doi.org/10.1101/gad.190678.112>.
- [19] Friend SH, Bernards R, Rogelj S, Weinberg RA, Rapaport JM, Albert DM, et al. 1986. A human DNA segment with properties of the gene that predisposes to retinoblastoma and osteosarcoma. *Nature* 323:643-646. <https://doi.org/10.1038/323643a0>.
- [20] Galluzzi L, Pietrocola F, Bravo-San Pedro JM, Amaravadi RK, Baehrecke EH, Cecconi F, et al. 2015. Autophagy in malignant transformation and cancer progression. *EMBO J.* 34:856-880. <https://doi.org/10.15252/emboj.201490784>.
- [21] Garrison SP, Jeffers JR, Yang C, Nilsson JA, Hall MA, Rehg JE, et al. 2008. Selection against PUMA gene expression in Myc-driven B-cell lymphomagenesis. *Mol. Cell Biol.* 28:5391-5402. <https://doi.org/10.1128/MCB.00907-07>.
- [22] Gewirtz DA. 2013. Autophagy and senescence: a partnership in search of definition. *Autophagy* 9:808-812. <https://doi.org/10.4161/aut.23922>.
- [23] Gole B, Wiesmuller L. 2015. Leukemogenic rearrangements at the mixed lineage leukemia gene (MLL)-multiple rather than a single mechanism. *Front. Cell Dev. Biol.* 3:41. <https://doi.org/10.3389/fcell.2015.00041>.
- [24] Gregory CD, Pound JD. 2011. Cell death in the neighbourhood: direct microenvironmental effects of apoptosis in normal and neoplastic tissues. *J. Pathol.* 223:177-194. <https://doi.org/10.1002/path.2792>.
- [25] Grivennikov SI, Greten FR, Karin M. 2010. Immunity, inflammation, and cancer. *Cell* 140:883-899. <https://doi.org/10.1016/j.cell.2010.01.025>.
- [26] Gupta A, Yang Q, Pandita RK, Hunt CR, Xiang T, Misri S, et al. 2009. Cell cycle checkpoint defects contribute to genomic instability in PTEN deficient cells independent of DNA DSB repair. *Cell Cycle* 8:2198-2210. <https://doi.org/10.4161/cc.8.14.8947>.
- [27] Hardwick JM, Soane L. 2013. Multiple functions of BCL-2 family proteins. *Cold Spring Harb. Perspect. Biol.* 5:a008722. <https://doi.org/10.1101/cshperspect.a008722>.
- [28] Hars ES, Lyu YL, Lin CP, Liu LF. 2006. Role of apoptotic nuclease caspase-activated DNase in etoposide-induced treatment-related acute myelogenous leukemia. *Cancer Res.* 66:8975-8979. <https://doi.org/10.1158/0008-5472.CAN-06-1724>.
- [29] Huang Q, Fang Li, Xinjian Liu, Wenrong Li, Wei Shi, Fei-Fei Liu, et al. 2011. Caspase 3-mediated stimulation of tumor cell repopulation during cancer radiotherapy. *Nat. Med.* 17:860-866. <https://doi.org/10.1038/nm.2385>.
- [30] Huh JR, Guo M, Hay BA. 2004. Compensatory proliferation induced by cell death in the Drosophila wing disc requires activity of the apical cell death caspase Dronc in a nonapoptotic role. *Curr. Biol.* 14:1262-1266. <https://doi.org/10.1016/j.cub.2004.06.015>.
- [31] Hyman BT, Yuan J. 2012. Apoptotic and non-apoptotic roles of caspases in neuronal physiology and pathophysiology. *Nat. Rev. Neurosci.* 13:395-406. <https://doi.org/10.1038/nrn3228>.
- [32] Ichim G, Lopez J, Ahmed SU, Muthalagund N, Giampazolias E, Delgado ME, et al. 2015. Limited mitochondrial permeabilization causes DNA damage and genomic instability in the absence of cell death. *Mol. Cell* 57:860-872. <https://doi.org/10.1016/j.molcel.2015.01.018>.
- [33] Inami Y, Waguri S, Sakamoto A, Kouno T, Nakada K, Hino O, et al. 2011. Persistent activation of Nrf2 through p62 in hepatocellular carcinoma cells. *J. Cell Biol.* 193:275-284. <https://doi.org/10.1083/jcb.201102031>.
- [34] Irby RB, Mao W, Coppola D, Kang J, Loubeau JM, Trudeau W, et al. 1999. Activating SRC mutation in a subset of advanced human colon cancers. *Nat. Genet.* 21:187-190. <https://doi.org/10.1038/5971>.
- [35] Islam MT. 2016a. Membrane marker sensory strategy (MMSS) is a new concept in cancer therapy: A hypothesis. *Int. J. Pharm. Pharm. Sci.* 8(12):314-317.
- [36] Islam MT. 2016b. Crucial challenges in epigenetic cancer therapeutic strategy yet to be resolved. *Int. J. Pharm. Pharm. Sci.* 8(12):1-6. <https://doi.org/10.22159/ijpps.2016v8i12.14510>.
- [37] Jeffers JR, Parganas E, Lee Y, Yang C, Wang J, Brennan J, et al. 2003. Puma is an essential mediator of p53-dependent and -independent apoptotic pathways. *Cancer Cell* 4:321-328. [https://doi.org/10.1016/S1535-6108\(03\)00244-7](https://doi.org/10.1016/S1535-6108(03)00244-7).
- [38] Keniry M, Parsons R. 2008. The role of PTEN signaling perturbations in cancer and in targeted therapy. *Oncogene* 27:5477-5485. <https://doi.org/10.1038/ncr.2008.248>.
- [39] Kenzelmann Broz D, Attardi LD. 2010. In vivo analysis of p53 tumor suppressor function using genetically engineered mouse models. *Carcinogen* 31:1311-1318. <https://doi.org/10.1093/carcin/bgp331>.
- [40] Kilbride SM, Prehn JH. 2013. Central roles of apoptotic proteins in mitochondrial function. *Oncogene* 32:2703-2711. <https://doi.org/10.1038/ncr.2012.348>.
- [41] Kim M-J, Woo S-J, Yoon C-H, Lee J-S, An S, Choi Y-H, et al. 2011. Involvement of autophagy in oncogenic K-ras-induced malignant cell transformation. *J. Biol. Chem.* 286:12924-12932. <https://doi.org/10.1074/jbc.M110.138958>.
- [42] Kimmelman AC. 2011. The dynamic nature of autophagy in cancer. *Genes Dev.* 25:1999-2010. <https://doi.org/10.1101/gad.17558811>.

- [43] Kimmelman AC. 2015. Metabolic dependencies in RAS-driven cancers. *Clin. Cancer Res.* 21:1828-1834. <https://doi.org/10.1158/1078-0432.CCR-14-2425>.
- [44] Kirkin V, McEwan DG, Novak I, Dikic I. 2009. A role for ubiquitin in selective autophagy. *Mol. Cell* 34:259-269. <https://doi.org/10.1016/j.molcel.2009.04.026>.
- [45] Kruiswijk F, Labuschagne CF, Voudsen KH. 2015. P53 in survival, death and metabolic health: a life guard with a licence to kill. *Nat. Rev. Mol. Cell Biol.* 16:393-405. <https://doi.org/10.1038/nrm4007>.
- [46] Kurtova AV, Xiao J, Mo Q, Pazhanisamy S, Krasnow R, Lerner SP, et al. 2015. Blocking PGE<sub>2</sub>-induced tumour repopulation abrogates bladder cancer chemoresistance. *Nature* 517:209-213. <https://doi.org/10.1038/nature14034>.
- [47] Lain S, Hollick JJ, Campbell J, Staples OD, Higgins M, Aoubala M, et al. 2008. Discovery, in vivo activity, and mechanism of action of a small-molecule p53 activator. *Cancer Cell* 13:454-463. <https://doi.org/10.1016/j.ccr.2008.03.004>.
- [48] Lamfers ML, Grill J, Dirven CM, Van Beusechem VW, Geogerger B, Van Den Berg J, et al. 2002. Potential of the conditionally replicative adenovirus Ad5-Delta24RGD in the treatment of malignant gliomas and its enhanced effect with radiotherapy. *Cancer Res.* 62:5736-5742.
- [49] Lee MN, Tseng RC, Hsu HS, Chen JY, Tzao C, Ho WL, et al. 2007. Epigenetic inactivation of the chromosomal stability control genes BRCA1, BRCA2, and XRCC5 in non-small cell lung cancer. *Clin. Cancer Res.* 13:832-838. <https://doi.org/10.1158/1078-0432.CCR-05-2694>.
- [50] Leffers N, Lambeck AJ, Gooden MJ, Hoogeboom BN, Wolf R, Hamming IE, et al. 2009. Immunization with a P53 synthetic long peptide vaccine induces P53-specific immune responses in ovarian cancer patients, a phase II trial. *Int. J. Cancer* 125:2104-2113. <https://doi.org/10.1002/ijc.24597>.
- [51] Letai AG. 2008. Diagnosing and exploiting cancer's addiction to blocks in apoptosis. *Nat. Rev. Cancer* 8:121-132. <https://doi.org/10.1038/nrc2297>.
- [52] Li F, Huang Q, Chen J, Peng Y, Roop D, Bedford JS, et al. 2010. Apoptotic cells activate the "phoenix rising" pathway to promote wound healing and tissue regeneration. *Sci. Signal* 3:ra13. <https://doi.org/10.1126/scisignal.2000634>.
- [53] Liu X, He Y, Li F, Huang Q, Kato TA, Hall RP, et al. 2015. Caspase-3 promotes genetic instability and carcinogenesis. *Mol. Cell* 58:284-296. <https://doi.org/10.1016/j.molcel.2015.03.003>.
- [54] Lock R, Roy S, Kenific CM, Su JS, Salas E, Ronen SM, et al. 2011. Autophagy facilitates glycolysis during ras-mediated oncogenic transformation. *Mol. Biol. Cell* 22:165-178. <https://doi.org/10.1091/mbc.E10-06-0500>.
- [55] Lovric MM, Hawkins CJ. 2010. TRAIL treatment provokes mutations in surviving cells. *Oncogene* 29:5048-5060. <https://doi.org/10.1038/onc.2010.242>.
- [56] Luke JJ, Van De Wetering CI, Knudson CM. 2003. Lymphoma development in Bax transgenic mice is inhibited by Bcl-2 and associated with chromosomal instability. *Cell Death Differ.* 10:740-748. <https://doi.org/10.1038/sj.cdd.4401233>.
- [57] Manning BD, Cantley LC. 2007. AKT/PKB signaling: navigating downstream. *Cell* 129:1261-1274. <https://doi.org/10.1016/j.cell.2007.06.009>.
- [58] Martin SA, McCabe N, Mullarkey M, Cummins R, Burgess DJ, Nakabeppu Y, et al. 2010. DNA polymerases as potential therapeutic targets for cancers deficient in the DNA mismatch repair proteins MSH2 or MLH1. *Cancer Cell* 17:235-248. <https://doi.org/10.1016/j.ccr.2009.12.046>.
- [59] Marusyk A, Porter CC, Zaberezhnyy V, DeGregori J. 2010. Irradiation selects for p53-deficient hematopoietic progenitors. *PLoS Biol.* 8:e1000324. <https://doi.org/10.1371/journal.pbio.1000324>.
- [60] Mathew R, Karp CM, Beaudoin B, Vuong N, Chen G, Chen H-Y, et al. 2009. Autophagy suppresses tumorigenesis through elimination of p62. *Cell* 137:1062-1075. <https://doi.org/10.1016/j.cell.2009.03.048>.
- [61] Mattson MP. 2000. Apoptosis in neurodegenerative disorders. *Nat. Rev. Mol. Cell Biol.* 1:120-129. <https://doi.org/10.1038/35040009>.
- [62] Michaelis M, Rothweiler F, Barth S, Cinatl J, van Rikxoort M, Löschmann N, et al. 2011. Adaptation of cancer cells from different entities to the MDM2 inhibitor nutlin-3 results in the emergence of p53-mutated multi-drug-resistant cancer cells. *Cell Death Dis.* 2:e243. <https://doi.org/10.1038/cddis.2011.129>.
- [63] Michalak EM, Jansen ES, Happo L, Cragg MS, Tai L, Smyth GK, et al. 2009. Puma and to a lesser extent Noxa are suppressors of Myc-induced lymphomagenesis. *Cell Death Differ.* 16:684-696. <https://doi.org/10.1038/cdd.2008.195>.
- [64] Muller FL, Colla S, Aquilanti E, Manzo VE, Genovese G, Lee J, et al. 2012. Passenger deletions generate therapeutic vulnerabilities in cancer. *Nature* 488:337-342. <https://doi.org/10.1038/nature11331>.
- [65] Mutka SC, Yang WQ, Dong SD, Ward SL, Craig DA, Timmermans PB, et al. 2009. Identification of nuclear export inhibitors with potent anticancer activity in vivo. *Cancer Res.* 69:510-517. <https://doi.org/10.1158/0008-5472.CAN-08-0858>.
- [66] Nagata S. 2010. Apoptosis and autoimmune diseases. *Ann. NY Acad. Sci.* 1209:10-16. <https://doi.org/10.1111/j.1749-6632.2010.05749.x>.
- [67] Nemunaitis J, Ganly I, Khuri F, Arseneau J, Kuhn J, McCarty T, et al. 2010. Selective replication and oncolysis in p53 mutant tumors with ONYX-015, an E1B-55kD gene-deleted adenovirus, in patients with advanced head and neck cancer: a phase II trial. *Cancer Res.* 60:6359-6366.
- [68] Noy R, Pollard JW. 2014. Tumor-associated macrophages: from mechanisms to therapy. *Immunity* 41:49-61. <https://doi.org/10.1016/j.immuni.2014.06.010>.
- [69] Pedro JM, Wei Y, Sica V, Maiuri MC, Zou Z, Kroemer G, Levine B. 2015. BAX and BAK1 are dispensable for ABT-737-induced dissociation of the BCL2-BECN1 complex and autophagy. *Autophagy* 11:452-459. <https://doi.org/10.1080/15548627.2015.1017191>.
- [70] Perez-Garijo A, Martin FA, Morata G. 2004. Caspase inhibition during apoptosis causes abnormal signalling and developmental aberrations in *Drosophila*. *Development* 131:5591-5598. <https://doi.org/10.1242/dev.01432>.
- [71] Pérez-Mancera PA, Young ARJ, Narita M. 2014. Inside and out: the activities of senescence in cancer. *Nat. Rev. Cancer* 14:547-558. <https://doi.org/10.1038/nrc3773>.
- [72] Peter ME, Hadji A, Murmann AE, Brockway S, Putzbach W, Patanayak A, et al. 2015. The role of CD95 and CD95 ligand in cancer. *Cell Death Differ.* 22:885-886. <https://doi.org/10.1038/cdd.2015.25>.
- [73] Puc J, Keniry M, Li HS, Pandita TK, Choudhury AD, Memeo L, et al. 2005. Lack of PTEN sequesters CHK1 and initiates genetic instability. *Cancer Cell* 7:193-204. <https://doi.org/10.1016/j.ccr.2005.01.009>.
- [74] Qu X, Yu J, Bhagat G, Furuya N, Hibshoosh H, Troxel A, et al. 2003. Promotion of tumorigenesis by heterozygous disruption of the beclin 1 autophagy gene. *J. Clin. Invest.* 112:1809-1820. <https://doi.org/10.1172/JCI20039>.
- [75] Quon KC, Berns A. 2001. Haplo-insufficiency? Let me count the ways. *Genes Dev.* 15:2917-2921. <https://doi.org/10.1101/gad.949001>.
- [76] Rajasagi M, Shukla SA, Fritsch EF, Keskin DB, DeLuca D, Carmona E, et al. 2014. Systematic identification of personal tumor-specific neoantigens in chronic lymphocytic leukemia. *Blood* 124:453-462. <https://doi.org/10.1182/blood-2014-04-567933>.
- [77] Rao RC, Dou Y. 2015. Hijacked in cancer: the KMT2 (MLL) family of methyltransferases. *Nat. Rev. Cancer* 15:334-346. <https://doi.org/10.1038/nrc3929>.
- [78] Rehm M, Huber HJ, Dussmann H, Prehn JH. 2006. Systems analysis of effector caspase activation and its control by X-linked inhibitor of apoptosis protein. *EMBO J.* 25:4338-4349. <https://doi.org/10.1038/sj.emboj.7601295>.
- [79] Ryo H, Gorenc T, Steller H. 2004. Apoptotic cells can induce compensatory cell proliferation through the JNK and the Wingless signaling pathways. *Dev. Cell* 7:491-501. <https://doi.org/10.1016/j.devcel.2004.08.019>.
- [80] Sandilands E, Serrels B, Wilkinson S, Frame MC. 2012. Src-dependent autophagic degradation of Ret in FAK-signalling-defective cancer cells. *EMBO Rep.* 13:733-740. <https://doi.org/10.1038/embor.2012.92>.
- [81] Sandilands E, Serrels S, McEwan DG, Morton JP, Macagno JP, McLeod K, et al. 2011. Autophagic targeting of Src promotes cancer cell survival following reduced FAK signaling. *Nat. Cell Biol.* 14:51-60. <https://doi.org/10.1038/ncb2386>.
- [82] Schuler M, Rochlitz C, Horowitz JA, Schlegel J, Perruchoud AP, Kommos F, et al. 1998. A phase I study of adenovirus-mediated wild-type p53 gene transfer in patients with advanced non-small cell lung cancer. *Hum. Gene Ther.* 9:2075-2082. <https://doi.org/10.1089/hum.1998.9.14-2075>.
- [83] Schuler PJ, Harasymczuk M, Visus C, Deleo A, Trivedi S, Lei Y, et al. 2014. Phase I dendritic cell p53 peptide vaccine for head and neck cancer. *Clin. Cancer Res.* 20:2433-2444. <https://doi.org/10.1158/1078-0432.CCR-13-2617>.
- [84] Seshadri R, Kutlaca RJ, Trainor K, Matthews C, Morley AA. 1987. Mutation rate of normal and malignant human lymphocytes. *Cancer Res.* 47:407-409.

- [85] Shall S, Gaymes T, Farzaneh F, Curtin N, Mufti GJ. 2011. The use of PARP inhibitors in cancer therapy: use as adjuvant with chemotherapy or radiotherapy; use as a single agent in susceptible patients; techniques used to identify susceptible patients. *Methods Mol. Biol.* 780:239-266. [https://doi.org/10.1007/978-1-61779-270-0\\_15](https://doi.org/10.1007/978-1-61779-270-0_15).
- [86] Somasekharan SP, Koc M, Morizot A, Micheau O, Sorensen PH, Gaide O, et al. 2013. TRAIL promotes membrane blebbing, detachment and migration of cells displaying a dysfunctional intrinsic pathway of apoptosis. *Apoptosis* 18:324-336. <https://doi.org/10.1007/s10495-012-0782-6>.
- [87] Spychala J. 2000. Tumor-promoting functions of adenosine. *Pharmacol. Ther.* 87:161-173. [https://doi.org/10.1016/S0163-7258\(00\)00053-X](https://doi.org/10.1016/S0163-7258(00)00053-X).
- [88] Stanford JC, Young C, Hicks D, Owens P, Williams A, Vaught DB, et al. 2014. Efferocytosis produces a prometastatic landscape during postpartum mammary gland involution. *J. Clin. Invest.* 124:4737-4752. <https://doi.org/10.1172/JCI76375>.
- [89] Sulzmaier FJ, Jean C, Schlaepfer DD. 2014. FAK in cancer: mechanistic findings and clinical applications. *Nat. Rev. Cancer* 14:598-610. <https://doi.org/10.1038/nrc3792>.
- [90] Sun N, Yun J, Liu J, Malide D, Liu C, Rovira II, et al. 2015. Measuring in vivo mitophagy. *Mol. Cell* 60:685-696. <https://doi.org/10.1016/j.molcel.2015.10.009>.
- [91] Swisher SG, Roth JA, Nemunaitis J, Lawrence DD, Kemp BL, Carrasco CH, et al. 1999. Adenovirus-mediated p53 gene transfer in advanced non-small-cell lung cancer. *J. Natl. Cancer Inst.* 91:763-771. <https://doi.org/10.1093/jnci/91.9.763>.
- [92] Tang J, Chen Y, Cui R, Li D, Xiao L, Lin P, et al. 2015. Upregulation of fractalkine contributes to the proliferative response of prostate cancer cells to hypoxia via promoting the G1/S phase transition. *Mol. Med. Rep.* 12:7907-7914. <https://doi.org/10.3892/mmr.2015.4438>.
- [93] Tao JJ, Castel P, Radosevic-Robin N, Elkabets M, Auricchio N, Aceto N, et al. 2014. Antagonism of EGFR and HER3 enhances the response to inhibitors of the PI3K-Akt pathway in triple-negative breast cancer. *Sci. Signal* 7:ra29. <https://doi.org/10.1126/scisignal.2005125>.
- [94] Tardaguila M, Manes S. 2013. CX3CL1 at the crossroad of EGF signals: relevance for the progression of ERBB2 breast carcinoma. *Oncotarget* 4:25669. <https://doi.org/10.4161/onci.25669>.
- [95] Thomas SM, Brugge JS. 1997. Cellular functions regulated by Src family kinases. *Ann. Rev. Cell Dev. Biol.* 13:513-609. <https://doi.org/10.1146/annurev.cellbio.13.1.513>.
- [96] Thorburn A. 2014. Autophagy and its effects: making sense of double-edged swords. *PLoS Biol.* 12:e1001967. <https://doi.org/10.1371/journal.pbio.1001967>.
- [97] Trinchieri G. 2012. Cancer and inflammation: an old intuition with rapidly evolving new concepts. *Ann. Rev. Immunol.* 30:677-706. <https://doi.org/10.1146/annurev-immunol-020711-075008>.
- [98] Vakkila J, Lotze MT. 2004. Inflammation and necrosis promote tumour growth. *Nat. Rev. Immunol.* 4:641-648. <https://doi.org/10.1038/nri1415>.
- [99] Villunger A, Michalak EM, Coultas L, Müllauer F, Böck G, Ausserlechner MJ, et al. 2003. p53-and drug-induced apoptotic responses mediated by BH3-only proteins Puma and Noxa. *Sci.* 302:1036-1038. <https://doi.org/10.1126/science.1090072>.
- [100] Vodenicharov MD, Sallmann FR, Satoh MS, Poirier GG. 2000. Base excision repair is efficient in cells lacking poly (ADP-ribose) polymerase1. *Nucleic Acids Res.* 28:3887-3896. <https://doi.org/10.1093/nar/28.20.3887>.
- [101] Von Karstedt S, Conti A, Nobis M, Montinaro A, Hartwig T, Lemke J, et al. 2015. Cancer cell-autonomous TRAIL-R signaling promotes KRAS-driven cancer progression, invasion, and metastasis. *Cancer Cell* 27:561-573. <https://doi.org/10.1016/j.ccell.2015.02.014>.
- [102] Wang Y, Suh YA, Fuller MY, Jackson JG, Xiong S, Terzian T, et al. 2011. Restoring expression of wild-type p53 suppresses tumor growth but does not cause tumor regression in mice with a p53 missense mutation. *J. Clin. Invest.* 121:893-904. <https://doi.org/10.1172/JCI44504>.
- [103] Weissmueller S, Machado E, Saborowski M, Morris JP 4th, Wagenblast E, Davis CA, et al. 2014. Mutant p53 drives pancreatic cancer metastasis through cell-autonomous PDGF receptor beta signaling. *Cell* 157:382-394. <https://doi.org/10.1016/j.cell.2014.01.066>.
- [104] Weston VJ, Oldreive CE, Skowronska A, Oscier DG, Pratt G, Dyer MJ, et al. 2010. The PARP inhibitor olaparib induces significant killing of ATM-deficient lymphoid tumor cells *in vitro* and *in vivo*. *Blood* 116:4578-4587. <https://doi.org/10.1182/blood-2010-01-265769>.
- [105] White E. 2015. The role for autophagy in cancer. *J. Clin. Invest.* 125:42-46. <https://doi.org/10.1172/JCI73941>.
- [106] Zelenay S, van der Veen AG, Böttcher JP, Snelgrove KJ, Rogers N, Acton SE, et al. 2015. Cyclooxygenase-dependent tumor growth through evasion of immunity. *Cell* 162:1257-1270. <https://doi.org/10.1016/j.cell.2015.08.015>.
- [107] Zhang H, Baehrecke EH. 2015. Eaten alive: novel insights into autophagy from multicellular model systems. *Trends Cell Biol.* 25:376-387. <https://doi.org/10.1016/j.tcb.2015.03.001>.