



Double right renal vein: clinical and surgical implications and review of the literature

José Aderval Aragão ^{1*}, Helen Lima Gomes ², Hiago Vinícius Dantas Costa ³,
Israel Santos Marcelo ³, Paula Santos Nunes ⁴

¹ *Associado Professor I, Department of Morphology and the Postgraduate Physical Education and Applied Health Sciences Programs, Federal University of Sergipe (UFS), and Titular Professor of the Medical School, Tiradentes University (UNIT), Aracaju, Sergipe, Brazil*

² *Medical Student, Tiradentes University (UNIT), Aracaju, Sergipe, Brazil*

³ *Medical Student, Federal University of Sergipe (UFS), Aracaju, Sergipe, Brazil*

⁴ *Adjunct Professor I, Department of Morphology of Federal University of Sergipe (UFS), Aracaju, Sergipe, Brazil*
*Corresponding author E-mail: jaafelipe@infonet.com.br

Copyright © 2015 José Aderval Aragão et al. This is an open access article distributed under the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Context: Variations in renal vascular morphology are relatively common and involve both the renal arteries and the renal veins. Presence of supernumerary vessels is the variation most frequently encountered. Knowledge of such findings is valuable for urologists with regard to kidney transplantation, nephrectomy, vascular anastomosis, selective catheterization and many other surgical procedures on the kidneys.

Objective: To report on a case of duplication of the right renal vein and its clinical and surgical implications.

Case report: In one dissected specimen from a male cadaver, the presence of two right renal veins parallel to each other was observed in a position anteroinferior to the right renal artery. One vein emerged from the upper portion of the renal hilum, while the other emerged from the lower portion of the hilum. The two veins drained separately into the inferior vena cava, and neither of them received any tributaries.

Conclusion: Knowledge of these vascular variations is important for urologists, vascular surgeons and radiologists, given that performing angiography prior to surgical interventions in the retroperitoneal space avoids complications, especially with regard to kidney transplantation.

Keywords: *Anatomical Variations; Renal Veins; Inferior Vena Cava; Kidney Transplantation; Accessory Renal Vessels.*

1. Introduction

The renal veins occur bilaterally and drain each of the kidneys into the inferior vena cava. Because the right kidney is closer to the inferior vena cava, the right renal vein is generally shorter and rarely receives tributaries. On the other hand, the left renal vein, which is longer, may receive several tributaries [1], [2].

Variations in the origin, path and termination of the renal veins, along with vascular malformations that are found in the renal hilum involving both arteries and veins, result mainly from errors of embryonic development [3], [4].

Several vascular anatomical variations in the renal hilum have been studied. The appearance of supernumerary vessels is the variation most frequently encountered among the renal morphological variations [1], [2], [5-8].

Knowledge of these variations, along with any other variations, is clinically important, because it allows surgeons to make better preoperative evaluations and thus minimize the risk of surgical complications [9]. The present study had the aim of reporting on a case of duplication of the right renal vein and the clinical and surgical implications of this anatomical variation.

2. Case report

In one dissected specimen from a male cadaver of dark skin color, with apparent age between 50 and 55 years, belonging to the Anatomy Laboratory of Tiradentes University, the presence of two right renal veins parallel to each other was observed in a position anteroinferior to the renal artery. These two veins drained separately into the inferior vena cava without receiving tributaries. The first of these veins emerged from the upper third of the renal hilum, close to the entry point of the right renal artery, and the second of these veins emerged from the lower third of the hilum (Fig. 1). On emerging from the renal hilum, the upper renal vein (which was of larger diameter) took an oblique anteroinferior direction, underwent a constriction followed by dilatation, probably due to slight compression of one of the segmental renal arteries, and emptied into the inferior vena cava, forming an acute angle with this (Fig. 2). Meanwhile, on emerging from the renal hilum, the inferior renal vein (which was smaller in diameter) took an ascending direction and then made a curve of approximately 120° downwards and emptied into the inferior vena cava, forming approximately a right angle at a posteroinferior position in relation to the upper renal vein (Fig. 1,2).




Fig. 1: Double Renal Vein Draining Into the Inferior Vena Cava




Fig. 2: Direction of the Double Renal Veins, from Their Emergence from the Renal Hilum to Their Outlet in the Inferior Vena Cava

3. Discussion

Clear knowledge of the embryology of the renal vein and inferior vena cava system is very important given its high complexity, and remains the best way to understand the variations and anomalies of this system. The retroperitoneal venous system develops from three parallel fetal venous systems: posterior cardinal, subcardinal and supracardinal. Anastomosis between the subcardinal and supracardinal systems forms the renal vein segment, and the supracardinal system gives rise to the postrenal segment of the inferior vena cava, below the diaphragm. This receives two renal veins from each side (one ventral and the other dorsal) and, during the course of the 8th week of fetal life. The two veins merge to form a single vein on each side. An error of fusion of these two vessels, thus causing their persistence, may result in bilateral or unilateral duplication of these vessels [10].

Occurrences of multiple renal veins are relatively common, and several authors have reported large numbers of venous variations resulting from embryonic developmental errors [1], [5], [11-13]. According to Kaneko, Kobayashi and Okada [14], the renal veins present fewer variations than do the renal arteries, and multiple renal veins are more common on the right side than on the left-side [8].

In our study, multiplicity of the renal veins was found on the right side. One emerged from the upper third and the other from the lower third of the renal hilum. The upper renal vein emptied into the inferior vena cava forming an acute angle, while the lower renal vein formed a right angle with the inferior vena cava. Similar findings were reported by Verma et al [6].

According to Satyapal [5], this angle was usually $60^\circ \pm 17^\circ$ on the right side. This author stated that large variations in this measurement would not be surprising, given that the kidneys are usually regarded as "floating viscera" that vary in the position with posture and with respiratory movements. Moreover, in vivo measurements differ from those made in cadavers.

In the present study, both of the renal veins emptied into the inferior vena cava without having received any tributaries, and this has also been reported by other authors [1], [15-17]. In most cases, these variations of duplication of the renal veins are clinically silent and remain unnoticed until they are discovered in the course of venography, operations or

autopsy [18]. For kidney transplantation surgeons, the morphology of the renal vein takes on special significance, given that its variations influence the technical viability of the operation and may restrict mobilization of the vein [7,19]. Nonetheless. Some anatomic variations in the renal veins may be correlated with clinical symptoms, as is the case of pain in the pelvic varices or varicocele. Furthermore, this multiplicity is more common in women than in men, as demonstrated by Arslan et al., Karazincir et al. and Koc et al [20-22]., Therefore, description of the anatomic variations of the renal veins is necessary with regard to investigating the effects of these variations on hemodynamics and renal function, particularly in cases of surgical interventions in the retroperitoneal space [2], [23].

4. Conclusion

Detailed anatomical knowledge of duplication of the renal veins and their main anomalies is extremely important for radiologists, vascular surgeons and urologists, since it provides guidance that will serve to prevent complications that might occur both in endovascular procedures and in surgery involving retroperitoneal access, especially given the increasing frequency of kidney transplantation surgery.

Acknowledgements

We thank Eufrazio dos Santos Cardoso; anatomy laboratory technician in the Tiradentes University, for their support in preparing the cadavers, since without their collaboration, this work could not have been accomplished.

References

- [1] Baptista-Silva JC, Veríssimo MJ, Castro MJ, Câmara AL, Pestana JO. Anatomical study of the renal veins observed during 342 living-donor nephrectomies. *Sao Paulo Med J.* 1997 May-Jun; 115 (3):1456-9. <http://dx.doi.org/10.1590/S1516-31801997000300011>.
- [2] Mir NS, Ul Hassan A, Rangrez R et al. Bilateral Duplication of Renal Vessels: Anatomical, Medical and Surgical perspective. *Int J Health Sci (Qassim).* 2008 Jul;2 (2):179-85.
- [3] Mathews R, Smith PA, Fishman EK, Marshall FF. Anomalies of the inferior vena cava and renal veins: embryologic and surgical considerations. *Urology.* 1999 May; 53 (5):873-80. [http://dx.doi.org/10.1016/S0090-4295\(99\)00007-2](http://dx.doi.org/10.1016/S0090-4295(99)00007-2).
- [4] Gillot C. La veine rénale gauche. *Anatomia Clinica.* 1978 avril; 1:135-55. <http://dx.doi.org/10.1007/BF01654493>.
- [5] Satyapal KS. The renal veins: A review. *Eur J Anat.* 2003 Jul; 7(S1):43-52.
- [6] Verma R, Kalra S, Rana K. Malformation of Renal and Testicular Veins– A case Report. *J Anat Soc India.* 2005; 54(1):29-31.
- [7] Krishnasamy N, Rao KGM, Somayaji SN, Koshy S, Rodrigues V. An unusual case of unilateral additional right renal artery and vein. *IJAV.* 2010; 3:9-11.
- [8] Gupta A, Gupta R, Singal R. Congenital Variations of Renal Veins: Embryological Background and Clinical Implications. *J Clin Diagn Res.* 2011 Nov; 5 (6):1140-43.
- [9] Toda R, Iguro Y, Moriyama Y, Hisashi Y, Masuda H, Sakata R. Double left renal vein associated with abdominal aortic aneurysm. *Ann Thorac Cardiovasc Surg.* 2001 Apr; 7 (2):113-5.
- [10] Biswas S, Chattopadhyay JC, Panicker H, Amblagan J, Ghosh SK. Variations in Renal and Testicular veins - A Case Report. *J. Anat. Soc. India.* 2006 Dec; 55(2):69-71.
- [11] Satyapal KS, Kalideen JM, Haffejee AA, Singh B, Robbs JV. Left renal vein variations. *Surg Radiol Anat.* 1999; 21 (1):77-81. <http://dx.doi.org/10.1007/BF01635058>.
- [12] Fernandes RMP, Conte FHP, Favorito LA, Abidu-Figueiredo M, Babinski MA. Triple right renal vein: An uncommon variation. *Int J Morphol.* 2005 Sep; 23 (3):231-33. <http://dx.doi.org/10.4067/S0717-95022005000300007>.
- [13] Bolla SR, Kumar S, Gorantla VR, Reddy S, Pamidi N, Jetti R. An unusual case of bilateral duplication of renal vessels: its clinical implications. *Int J Clin Surg Adv.* 2013 Jul; 1 (1):20-3.
- [14] Kaneko N, Kobayashi Y, Okada Y. Anatomic variations of the renal vessels pertinent to transperitoneal vascular control in the management of trauma. *Surgery.* 2008 May; 143 (5):616-22. <http://dx.doi.org/10.1016/j.surg.2008.02.003>.
- [15] Anson BJ, Cauldwell EW, Pick JW, Beaton LE. The anatomy of the para-renal system of veins, with comments on the renal arteries. *J Urol.* 1948 Nov; 60 (5):714-37.
- [16] Baptista-Silva JCC, Cal RGR, Castro MJ, Veríssimo MJM, Câmara ALG. Estudo anatômico das veias renais e suas variações, observação durante as nefrectomias em doadores vivos. *Rev Col Bras Cir* 1995; 22 (2):18.
- [17] Gesase AP. Rare origin of supernumerary renal vessels supplying the lower pole of the left kidney. *Ann Anat.* 2007; 189 (1):53-8. <http://dx.doi.org/10.1016/j.aanat.2006.06.012>.
- [18] Prakash C, Ali S, Khan SM. Anomalous left renal vein coursing behind aorta and draining into the left common iliac vein: A rare variant. *Int J Case Rep Images* 2014; 5 (8):553–557.
- [19] Senecaïl B, Bobeuf J, Forlodou P, Nonent M. Two rare anomalies of the left renal vein. *Surg Radiol Anat.* 2003 Nov-Dec; 25 (5-6):465-7 <http://dx.doi.org/10.1007/s00276-003-0164-4>.
- [20] Arslan H, Etlik O, Ceylan K, Temizoz O, Harman M, Kavan M. Incidence of retro-aortic left renal vein and its relationship with varicocele. *Eur Radiol.* 2005 Aug; 15 (8):1717-20. <http://dx.doi.org/10.1007/s00330-004-2563-2>.
- [21] Karazincir S, Balci A, Görtür S, Sumbas H, Kiper AN. Incidence of the retroaortic left renal vein in patients with varicocele. *J Ultrasound Med.* 2007 May; 26 (5):601-4.
- [22] Koc Z, Ulsan S, Oguzkurt L, Tokmak N. Venous variants and anomalies on routine abdominal multi-detector row CT. *Eur J Radiol.* 2007 Feb; 61 (2):267-78. <http://dx.doi.org/10.1016/j.ejrad.2006.09.008>.
- [23] Boyaci N, Karakas E, Dokumaci DS, Yildiz S, Cece H. Evaluation of left renal vein and inferior vena cava variations through routine abdominal multi-slice computed tomography. *Folia Morphol (Warsz).* 2014 May; 73 (2):159-63. <http://dx.doi.org/10.5603/FM.2014.0017>.

Legends of figures

Fig. 1: Double renal vein draining into the inferior vena cava

AA – Aortic artery

RRA – Right renal artery

LRA – Left renal artery

RK – Right kidney

SMA – Superior mesenteric artery

IMA – Inferior mesenteric artery

CA – Celiac artery

IVC – Inferior vena cava

RRV1 – Right renal vein 1

RRV2 – Right renal vein 2

RCIA – Right common iliac artery

LCIA – Left common iliac artery

RIIA – Right internal iliac artery

LIIA – Left internal iliac artery

REIA – Right external iliac artery

LEIA – Left external iliac artery

MSA – Median sacral artery

RU – Right ureter

Fig. 2: Direction of the double renal veins, from their emergence from the renal hilum to their outlet in the inferior vena cava

AA – Aortic artery

RRA – Right renal artery

LRA – Left renal artery

RK – Right kidney

SMA – Superior mesenteric artery

IMA – Inferior mesenteric artery

CA – Celiac artery

IVC – Inferior vena cava

RRV1 – Right renal vein 1

RRV2 – Right renal vein 2

RU – Right ureter

RSRA – Right segmental renal artery