



## Bilateral accessory renal arteries in a fetus: importance for surgical and radiological practice

José Aderval Aragão <sup>1\*</sup>, Danilo Primo Andrade Santana <sup>2</sup>, Ellen Caroline da Conceição Brandão <sup>2</sup>, Viviane Freitas Andrade <sup>3</sup>, Marianna Ribeiro de Menezes Freire <sup>4</sup>, Francisco Prado Reis <sup>5</sup>

<sup>1</sup> *Associado Professor I, Department of Morphology and the Postgraduate Physical Education and Applied Health Sciences Programs, Federal University of Sergipe (UFS), and Titular Professor of the Medical School, Tiradentes University (UNIT), Aracaju, Sergipe, Brazil*

<sup>2</sup> *Medical Student, Federal University of Sergipe (UFS), Aracaju, Sergipe, Brazil*

<sup>3</sup> *Postgraduate Student Applied Health Sciences Programs, Federal University of Sergipe (UFS), Aracaju, Sergipe, Brazil*

<sup>4</sup> *Medical Student, Tiradentes University (UNIT), Aracaju, Sergipe, Brazil*

<sup>5</sup> *Titular Professor, Medical School of Tiradentes University (UNIT), Aracaju, Sergipe, Brazil*

\*Corresponding author E-mail: jaafelipe@infonet.com.br

Copyright © 2015 José Aderval Aragão et al. This is an open access article distributed under the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Context:** Knowledge of anatomical variations of the renal vessels is considered important, in view of the increasing use of kidney transplantation, vascular reconstruction and imaging methods.

**Objective:** To report on a case of triple right renal artery, and double left renal artery and its clinical-surgical implications.

**Case report:** In a female fetal cadaver of age 30 weeks, multiple renal arteries were observed: triple in the right and double in the left renal artery. All the arteries were hilar and originated from the lateral face of the abdominal aorta. The lengths of the triple renal arteries were 17, 10 and 12 mm respectively for the upper, middle and lower arteries. The lengths of the double arteries were 10 and 12 mm respectively for the upper and lower arteries. There was no extrarenal segmentation.

**Conclusion:** Knowledge of possible variations in the renal arteries may be very useful for radiologists, urologists and vascular surgeons.

**Keywords:** Anatomy; Renal Artery; Urogenital Abnormalities; Kidney Transplantation.

## 1. Introduction

Anatomical knowledge of variations in the numbers of renal arteries is considered important, in view particularly of the gradually increasing use of invasive radiological procedures, comprising urological, vascular and kidney transplantation operations [1-5]. The present study has aim of reporting on a case of triple right renal artery, and double left renal artery in a human fetus.

## 2. Case report

After dissecting 60 kidneys from human fetuses in order to determine the frequency of multiple renal arteries, one case of triple and double renal arteries (right and left, respectively) was found in a female fetus of estimated age 30 weeks (hallux-calcaneus measurement (Fig. 1). All the arteries were hilar and originated from the lateral face of the abdominal aorta. The lengths of the triple renal arteries were 17, 10 and 12 mm respectively for the upper, middle and lower arteries, which were located at distances of 2, 3 and 5 mm from the superior mesenteric artery. Before entering the renal hilum, the upper and middle triple renal arteries originated two segmental renal arteries. The lengths of the double renal

arteries were 10 and 12 mm respectively for the upper and lower arteries, and were located 3 and 10 mm from the superior mesenteric artery. There was no extrarenal segmentation.




Fig. 1: Triple and Double Renal Arteries

### 3. Discussion

An additional renal artery, named variously as accessory, aberrant, anomalous or malformed artery, which is commonly presented in the literature, may arise from different portions of the aorta and end at different locations in the kidney. Although there is no exact differentiation between these terms, the latter ones should be applied when the structural changes are translated into dysfunction [6], [7].

According to authors such as Sykes [8], Willan, and Humpherson [9], the anatomical variations generally do not have any effect on body functions, although, they may influence the course of a disease, and also the clinical and surgical findings. The renal arteries have been the subject of many systematic studies [10], although the numbers of reports of cases of double and triple arteries have been much smaller [11]. Nevertheless, variations in the numbers of renal arteries are more common than variations in any other vessel [12], [13], and this can be explained by the ascension of the kidneys from the pelvis during embryonic development [14]. The frequency of variations may range from 8.7 to 75.6% [15], [16].

The frequency of occurrences of triple renal arteries has been found to range from 0.6 to 4.5% [1], [2], [4], [17-23]. In our study, the frequency of triple renal arteries was 3.33% among 60 kidneys dissected from 30 human fetuses. This number was similar to what was found by Busato Júnior and Ribas Filho [20], who evaluated 30 molds obtained from the kidneys of fresh human cadavers, which had been prepared by means of the injection-corrosion technique using acrylic copolymer, as used in dentistry. According to Merklin and Michels [17], Harrison et al [18], Sampaio and Passos [1], Gościcka et al [19], Busato Júnior, and Ribas, Filho [20], Khamanarong et al [4], Bordei [21], Bergman et al [22] and Kaneko et al [23], this frequency ranged from 0.6 to 1.9%, while according to Satyapal et al [2], it was 4.5%.

Knowledge of these anatomical variations in renal vascularization are of etiological, diagnostic, and therapeutic importance, both for surgeons and for radiologists [24], [25]. The presence of multiple renal arteries increases the complexity of surgical procedures such as kidney transplantation, implantation of vascular stents and vascular reconstruction, and also has clinical importance with regard to diagnosing renovascular hypertension [26].

### 4. Conclusion

Knowledge of the multiple variations in the renal hilum is of great importance in making diagnoses through imaging before surgical interventions, and also in reducing the surgical complications in kidney transplantations.

## References

- [1] Sampaio FJ, Passos MA. Renal arteries: anatomic study for surgical and radiological practice. *Surg Radiol Anat.* 1992; 14 (2):113-7. <http://dx.doi.org/10.1007/BF01794885>.
- [2] Satyapal KS, Haffeejee AA, Singh B, Ramsaroop L, Robbs JV, Kalideen JM. Additional renal arteries: incidence and morphometry. *Surg Radiol Anat.* 2001; 23 (1):33-8. <http://dx.doi.org/10.1007/s00276-001-0033-y>.
- [3] Baltacıoğlu F, Ekinçi G, Akpınar İN, Cımsıt NC, Tuğlular S, Akoğlu E. Endovascular treatment of renal artery stenosis: Technical and clinical results. *Tani Girisim Radyol.* 2003 Jun; 9(2):246-56.
- [4] Khamanarong K, Prachaney P, Utraravichien A, Tong-Un T, Sriparaya K. Anatomy of renal arterial supply. *Clin Anat.* 2004 May; 17(4):334-6. <http://dx.doi.org/10.1002/ca.10236>.
- [5] Petru B, Elena S, Dan I, Constantin D. The morphology and the surgical importance of the gonadal arteries originating from the renal artery. *Surg Radiol Anat.* 2007 Jul; 29(5):367-71. <http://dx.doi.org/10.1007/s00276-007-0224-2>.
- [6] Sa-udo JR, Vázquez R, Puerta J. Meaning and clinical interest of the anatomical variations in the 21st century. *Eur J Anat.* 2003 July; 7(Suppl 1):1-3.
- [7] Raikos A, Paraskevas GK, Natsis K, Tzikas A, Njau SN. Multiple variations in the branching pattern of the abdominal aorta. *Rom J Morphol Embryol.* 2010; 51(3):585-7.
- [8] Sykes D. The arterial supply of the human kidney with special reference to accessory renal arteries. *Br J Surg.* 1963 Jan; 50:368-74. <http://dx.doi.org/10.1002/bjs.18005022204>.
- [9] Willan PL, Humpherson JR. Concepts of variation and normality in morphology: important issues at risk of neglect in modern undergraduate medical courses. *Clin Anat.* 1999; 12(3):186-90. [http://dx.doi.org/10.1002/\(SICI\)1098-2353\(1999\)12:3<186::AID-CA7>3.0.CO;2-6](http://dx.doi.org/10.1002/(SICI)1098-2353(1999)12:3<186::AID-CA7>3.0.CO;2-6).
- [10] Aragão JA, de Oliveira Pacheco JM, Silva LA, Reis FP. Frequency of multiple renal arteries in human fetuses. *Surg Radiol Anat.* 2012 Mar; 34(2):133-6. <http://dx.doi.org/10.1007/s00276-011-0860-4>.
- [11] Olave E, Henríquez J, Puelma F, Cruzat C, Soto A. Arterias Renales Múltiples. *Int. J. Morphol.* 2007; 25(4):927-30. <http://dx.doi.org/10.4067/S01717-95022007000400040>.
- [12] Hollinshead, WH. The kidneys, ureters and suprarenal glands. In: *Anatomy for surgeons.* 2<sup>a</sup> Ed. New York USA, Harper & Row Publishers, 1971; 2:518-568.
- [13] Kuczera M, Gajda G, Gielecki JS. Digital analysis of the dynamics of the arterial supply to the human foetal kidneys. *Folia Morphol.* 2003 Jul; 62(4):381-384.
- [14] Moore KL, Persaud TVN. Período Fetal: da nona semana ao nascimento. In: *Embriologia clínica.* 8<sup>a</sup> ed. Rio de Janeiro: Elsevier, 2008; p. 104.
- [15] Fine H, Keen EN. The arteries of the human kidney. *J Anat.* 1966; 100:881-894.
- [16] Spanos PK, Simmons RL, Kjellstrand CM, Buselmeier TJ, Najarian JS. Kidney transplantation from living related donors with multiple vessels. *Am J Surg.* 1973; 125: 554 -558? [http://dx.doi.org/10.1016/0002-9610\(73\)90137-2](http://dx.doi.org/10.1016/0002-9610(73)90137-2).
- [17] Merklin RJ, Michels NA. The variant renal and suprarenal blood supply with data on the inferior phrenic, ureteral and gonadal arteries: a statistical analysis based on 185 dissections and review of the literature. *J Int Coll Surg.* 1958 Jan; 29(1, Part 1):41-76.
- [18] Harrison LH Jr, Flye MW, Seigler HF. Incidence of anatomical variants in renal vasculature in the presence of normal renal function. *Ann Surg.* 1978 Jul; 188(1):83-9. <http://dx.doi.org/10.1097/0000658-197807000-00014>.
- [19] Gościcka D, Szpinda M, Kochan J. Accessory renal arteries in human fetuses. *Ann Anat.* 1996 Dec; 178(6):559-63. [http://dx.doi.org/10.1016/S0940-9602\(96\)80118-X](http://dx.doi.org/10.1016/S0940-9602(96)80118-X).
- [20] Busato Júnior WFS, Ribas Filho JM. Estudo da Distribuição Arterial em Rins Humanos. *Arquivos Catarinenses de Medicina.* 2003 Jul-Sep; 32(3):21-27.
- [21] Bordei P, Sapte E, Ilescu D. Double renal arteries originating from the aorta. *Surg Radiol Anat.* 2004 Dec; 26(6):474-9. <http://dx.doi.org/10.1007/s00276-004-0272-9>.
- [22] Bergman RA, Afifi AK, Miyauchi R. Renal arteries: Illustrated encyclopedia of human anatomic variation: Opus II: Cardiovascular system: Arteries: Abdomen. 2005. Available from: <http://www.vh.org/adult/provider/anatomyvariants/cardiovascular/text/arteries/renal.html>.
- [23] Kaneko N, Kobayashi Y, Okada Y. Anatomic variations of the renal vessels pertinent to transperitoneal vascular control in the management of trauma. *Surgery.* 2008 May; 143(5):616-22. <http://dx.doi.org/10.1016/j.surg.2008.02.003>.
- [24] Çiçekbaşı AE, Ziyilan T, Salbacak A, Seker M, Büyükmumcu M, Tuncer I. An investigation of the origin, location and variations of the renal arteries in human fetuses and their clinical relevance. *Ann Anat.* 2005 Sep; 187(4):421-7. <http://dx.doi.org/10.1016/j.aanat.2005.04.011>.
- [25] Mazzucchi E, Sousa AA, Nahas WC, Antonopoulos IM, Piovesan AC, Arap S. Surgical complications after renal transplantation in grafts with multiple arteries. *Int. Braz J Urol.* 2005 Mar-Apr; 31(2): 125-130. <http://dx.doi.org/10.1590/S1677-55382005000200006>.
- [26] Pestemalci T, Mavi A, Yildiz YZ, Yildirim M, Gumusburun E. Bilateral triple renal arteries. *Saudi J Kidney Dis Transpl.* 2009 May; 20(3):468-70.

## Legends of figures

Figure 1 Triple and double renal arteries (RRA1, RRA2, RRA3 and LRA1, LRA2)

AO – Aorta

CA – Celiac artery

IMA - Inferior mesenteric artery

LK – Left kidney

LRA1 – Left renal artery1

LRA2 – Left renal artery2

LSG – Left suprarenal gland

RK – Right kidney

RRA1 – Right renal artery1

RRA2 – Right renal artery2

RRA3 – Right renal artery3

RSG – Right suprarenal gland

SMA – Superior mesenteric artery