

Allelopathic effects of *Thuja orientalis* L. and *Melia azedarach* L. on seed germination of *Penicillium americanum* L

Saeeda Bibi^{1*}, Rukhsana Jabeen², Asma Abdul Hayee², Saima Johar², Shehnaz Sher Ahmed², Shaista Hameed², Mahpara Fida Ahmed²

¹Department of Plant Sciences, Sardar Bahadur Khan Women's University, Quetta, Pakistan

²Department of Plant Sciences, SBK Women's University Quetta, Pakistan

*Corresponding author E-mail: saeeda_botany66@yahoo.com

Abstract

Aqueous extract of leaves, bark and seeds of *Melia azedarach* L. and *Thuja orientalis* L. were assayed at 1, 5 and 10g/L concentration with different time period to check their effect on seed germination, fresh and dry weight and seedling growth of *Penicillium americanum* L. Result revealed that aqueous extracts of both plants at all concentration and time period had significantly inhibited seed germination of *P. americanum* L. when compared with control. The inhibitory effect increases with increasing concentration of extracts and time period. The bark extract of all concentration of *T. orientalis* in 24hr and 48hrs show stimulation in seedling growth while other parts and extracts of all parts of *M. azedarach* in all concentration and time period show inhibition in all parameters. The order of inhibition when compared different parts of *M. azedarach* was seeds > bark > leaves while that of *T. orientalis* was leaves > bark > seeds. Hence it is concluded that aqueous extract of these plant contain water soluble allelochemicals which inhibit the seed germination and seedling growth of *P. americanum* L. It is suggested that these chemicals may be used as herbicides.

Keywords: *Melia azedarach* L.; *Thuja orientalis* L.; *Penicillium americanum* L.; Aqueous Extract; Seed Germination; Seedling Growth. Fresh and Dry Weight; Allelochemicals.

1. Introduction

Allelopathy can be defined as the ability of plant to stimulate or inhibit the growth of other plants in the environment by releasing chemicals or Allelopathy can also be defined as any direct or indirect, beneficial or harmful effects of one plant on other through the production of allelochemicals that it release into the environment [9]. Allelopathy is the relationship between plants by means of allelochemicals released into the environment and exists in natural plant community for long time period and it include both promotion and inhibition [13]. All parts of plants such as Leaves, Fruits, Stem, seeds and roots contain allelochemicals which are released into the environment by the process of volatilization, root exudation, leaching and decomposition [12]. These chemicals are known to affect development, reproduction, growth, germination and distribution of a number of plant species [14]. Germination and growth inhibitory effects of plants are associated to allelopathy. Allelopathy plays an important role in agro-ecosystem and natural ecosystem and has both stimulatory and inhibitory effect by releasing allelochemicals into the environment [3].

Melia azedarach L. is an evergreen tree which belongs to family Meliaceae. It is commonly known as chinaberry tree and closely related to neem. [7] It is one of most important specie because it contains variety of secondary metabolites such as triterpenoids and lemanoids. This plant is widely distributed and moderate sizes deciduous tree with cylindrical bark the leaves are opposite and alternate. This plant is widely used medicinally. [9]

Thuja orientalis belongs to family Cupressaceae. It is monoecious an evergreen tree. This plant contain of phyto constituents such as flavonoids and terpenoids that showed the biological activities [3]. It has great medicinal value. It has antiviral action and immune pharmacological potential [16].

2. Materials and methods

2.1. Extract preparation

One gram, 5g and 10g leaves, stems and seeds of *Melia azedarach* L. and *Juglanregia* L. were extracted in 100 ml of distilled water in separate Erlenmeyer flasks (250 ml) for 24, 48 and 72 h at room temperature. The extracts were filtered through ordinary filter papers. The extracts were stored in air tight glass vials and in a fridge at 4°C till further analysis.

2.2. Germination of seeds

The seeds were germinated on filter paper which was cut in round shape equal to diameter to Petri dish. Twice folded filter paper was placed at the base of Petri dish. Five seeds of *P. americanum* were placed randomly on filter paper. These Petri dishes were placed in a germinator at 20°C and 20% humidity. For all the treatments the biotic factors were same. Three replicates were used for each plant and for each extract. In controlled conditions only distilled water was used.

2.3. Measurements of parameters

To determine the allelopathic effects, seed germination, length of plumule and radical, moisture content of seedlings, fresh and dry weight of seedlings of *P. americanum* were noted against different concentrations of extracts. Readings were taken after 7 day. The length of plumule, radical was measured by scale. The fresh and dry weights of seedlings were taken by digital balance. The germination percentage was also calculated by following formula:

$$\text{Germination percentage} = \frac{\text{Number of germinated seeds}}{\text{Total number of seeds}} \times 100$$

3. Results

The present study was conducted to check the allelopathic effect of leaves, stem and seeds on germination rate, fresh and dry weight and seedling growth of *pennisitumamericanum*. The result were

3.1. Leaves extract of *thujaorientalis L.*

Dried leaves of 1gm, 5gm, 10gm were soaked for 24, 48, 72 hours. These extracts were used for seed treatments. The germination rate of control was 93%. The leaves extract in all concentration and time period show inhibition in germination rate. The fresh and dry weight of control was 0.2g and 0.07g. 1gm aqueous extract of leaf in 48hrs and (5, 10g) in 48hrs show stimulation in seedling growth and fresh and dry weight. (Table 1.1).

Table 1.1: Effect of Aqueous Leaf Extract of *Thujaorientalis* on Germination Rate, Fresh and Dry Weight.

S/n	Treatments	Germination percentage (%)	Fresh weight (g)	Dry weight (g)	Seedling growth Length of radical(cm)	Length of plumule(cm)
1.	Control	93	0.2	0.07	4.5 ± 0.6	3.19 ± 0.46
2.	1gm					
	24hours	40	0.07	0.03	3.2 ± 2.3	2.2 ± 0.7
	48hours	60	0.27	0.19	3.2 ± 2.2	2.1 ± 0.7
3.	5gm					
	24hours	50	0.06		1.4 ± 0.36	0.3 ± 0.2
	48hours	47	0.25	0.03	5.1 ± 2.5	3.3 ± 1.9
4.	10gm					
	24hours	47	0.06	0.03	2.4 ± 1.2	0.9 ± 0.4
	48hours	53	0.28	0.18	6.3 ± 1.8	4.7 ± 1.1
	72hours	67	0.03	0.13	4.3 ± 0.97	1.3 ± 0.2

3.2. Aqueous extract of stem

Aqueous extract of stem in all concentration show inhibition in germination percentage and other parameters but some extract 1g in (24hr, 72hr) and 5g in (48hr) show stimulation in seedling growth while other show inhibition in all parameters.(Table 1.2)

Table 1.2: Effect of Aqueous Extract of Stem of *T.Orientalis* on Seed Germination, Seedling Growth and Fresh and Dry Weight of *P. Americanum*.

S/n	Treatments	Germination percentage (%)	Fresh weight(gm)	Dry weight (gm)	Seedling growth Length of radical (cm)	Length of plumule (cm)
1	Control	93	0.2	0.07	4.5 ± 0.6	3.19 ± 0.46
2	1gm					
	24hours	53.3	0.14	0.033	6.618 ± 3.26	2.78 ± 2.280
	48hours	73	0.13	0.04	3.2 ± 1.4	2.3 ± 1.0
3	5gm					
	24hours	67	0.2	0.04	5.7 ± 4.5	0.5 ± 0.4
	48hours	60	0.14	0.04	4.1 ± 1.5	2.6 ± 0.3
4	10gm					
	24hours	67	0.22	0.04	5.7 ± 4.5	0.5 ± 0.4
	48hours	60	0.13	0.03	0.8 ± 0.3	0.5 ± 0.4
	72hours	30	0.05	0.01	3.9 ± 0.6	2.5 ± 0.3
					3.8 ± 0.7	1.9 ± 0.2
					1.8 ± 0.6	0.4 ± 0.2

3.3. Aqueous extracts of seed

Aqueous extracts of seed also show inhibition in germination rate and other parameters but some extract such as 5g and 1g in (24hr) show increase in fresh and dry weight while other extract show decrease in fresh and dry weight. 1g and 5 g in (24hr, 72hrs) also show stimulation in seedling growth. (Table 1.3)

Table 1.3: Effect of Aqueous Extract of Seeds of *T.Orientalis* on Seed Germination, Seedling Growth and Fresh and Dry Weight of *P. Americanum*.

S/n	Treatments	Germination percentage (%)	Fresh weight(gm)	Dry weight (gm)	Seedling growth Length of radical (cm)	Length of plumule (cm)
1	Control	93	0.2	0.07		
2	1gm					
	24hours	87	0.24	0.11	4.9 ± 1.1	6.5 ± 1.6
	48hours	53	0.14	0.03	8.0 ± 2.4	7.9 ± 5.1
3	5gm					
	24hours	40	0.09	0.04	2.6 ± 0.6	1.0 ± 0.5
	48hours	60	0.5	0.28	6.7 ± 0.6	9.01 ± 1.1
4	10gm					
	24hours	60	0.2	0.1	5.2 ± 0.5	8.8 ± 2.3
	48hours	40	0.08	0.02	2.5 ± 1.5	1.3 ± 0.6
	72hours	87	0.06	0.05	1.9 ± 0.9	1.2 ± 0.8
					2.5 ± 1.5	1.7 ± 0.6
					0.06 ± 0.00	0.06 ± 0.00

3.4. Meliaazedarch L.

3.4.1. Aqueous extract of leaves

The leaves extract of *M. azedarach* in all concentration in time period of 72h showed 0% germination.while in other time period showed inhibition in all parameters. (Table 1.4)

Table 1.4: Effect of Aqueous Extract of Leaves of *M. Azedarach* on Seed Germination, Seedling Growth and Fresh and Dry Weight of *P. Americanum*.

s/n	Treatments	Germination percentage	Fresh weight	Dry weight	Seedling growth	
		(%)	(g)	(g)	Length of radical	Length of plumule
1	Control	93	0.2	0.07	4.1 ± 1.7	1.6 ± 0.2
2	1gm					
	24hours	47	0.04	0.02	0.6 ± 0.4	0.13 ± 0.02
	48hours	20	0.02	0.01	1.1 ± 0.1	0.6 ± 0.4
3	5gm					
	24hours	40	0.05	0.01	0.2 ± 0.1	0.1 ± 0.1
	48hours	27	0.04	0.01	0.4 ± 0.00	0.8 ± 0.7
	72hours	0	0	0		
4	10gm					
	24hours	7	0.02	0.01	0.07 ± 0.03	0.2 ± 0.1
	48hours	20	0.02	0.005	0.1 ± 0.03	0.3 ± 0.1
	72hours	0	0	0		

3.4.2. Aqueous extract of stem

The aqueous extract of *M. azedarach* L in all concentration in time period of 72h showed 0% germination.while in other time period showed inhibition in all parameters.(Table 1.5)

Table1.5: Effect of Aqueous Extract of Stem of *M. Azedarach* on Seed Germination, Seedling Growth and Fresh and Dry Weight of *P. Americanum*.

s/n	Treatments	Germination percentage%	Fresh weight (g)	Dry weight (g)	Seedling growth	
					Length of radical(cm)	Length of plumule(cm)
1	Control	93	0.2	0.07	4.1 ± 1.7	1.6 ± 0.2
2	1gm					
	24	47	0.05	0.03	0.1+0.9	0.5 ± 0.3
	48	53.3	0.03	0.01	11.2 ± 1.1	0.4 ± 0.4
	72	53.3	0.1	0.1	1.7 ± 0.6	0.5 ± 0.5
3	5gm					
	24	60	0.08	0.04	0.3±0.4	0.8± 0.3
	48	47	0.1	0.03	2.9 ± 1.0	1.6±0.4
	72	40	0.06	0.03	0.8± 0.3	0.5± 0.4
4	10gm					
	24	47	0.1	0.03	0.6± 0.1	0.2±0.1
	48	87	0.1	0.04	3.0 ± 1.2	1.4 ± 0.9
	72	0	0	0		

3.4.3. Aqueous extracts of seeds

The aqueous extracts of seeds of *M. azedarach* in all concentrations and time period showed 0% germination.

4. Discussion

Allelopathy is the ability of plant to stimulate or inhibit the growth of plants by secreting many chemicals in the environment

[10].The allelopathic potential of aqueous extracts of different parts of both plants at different concentration and time period was evaluated on seed germination percentage, fresh and dry weight and seedling growth of *P.ammericanum*. All aqueous extract of *M. azedarach* markedly inhibited all parameters but inhibitory effect increases with increasing concentration and time period. Aqueous extract of seed had more inhibitory effect than other parts when compared. The chemicals which show allelopathic activity are present in different parts of plants including stem, leaves, flowers, seeds and fruits. These chemicals are released into the environment by means of leaching, root exudation, decomposition of residue and volatilization [8].

Aqueous extract of leaves and stem of *Calatropisprocera* had inhibitory effect on seedling growth, germination and fresh and dry weight of *Pennisetumamericanum* L. Both plants showed inhibition in seed germination. *Meliaazedarach* L. showed more inhibition than other plants. Seedling growth is not equally affected by all plants. *Thujaorientalis* L. in some concentration showed stimulation in seedling growth. Leaves and stem extract of *Rhazystricta* also inhibit germination rate and seedling growth of *P. americanum*L.leaves had more inhibitory effect than stem [5].The aqueous extract of seeds, leaves and stem of *Meliaazedarach* L.in all concentrations (1gm,5gm,10gm) showed inhibitory effect on all parameters of *Pennisetumamericanum* L which increases with increasing concentration of extract. The order of inhibition was seed> leaves > stem.

Alcoholic and aqueous extract of different parts (fruits, leaves and leaves wood mix) of *Meliaazedarach* L reduced seedling growth as well as germination rate of *Lectuca sativa*. Rate of inhibition varied with extract type and its concentration and order of inhibition was fruit extract> leaves extract> leaves and wood mix extract [7].

Aqueous extracts of seeds of *M. azedarach*L.showed complete inhibition in all concentrations. The percentage of germination is zero (0) in all concentrations.Inhibitory effect is due to water soluble allelochemicals. The aqueous extract of stem, leaves and seeds of *T.orientales* L reduced germination percentage and fresh and dry weight of *Pennisetumamericanum* L but show stimulation in seedling growth.

Leaves extract of *T. orientales* L in all concentrations and stem extract in 48hrs and 24hrs showed stimulation in seedling growth.The order of inhibition when compared to different parts of plant was leaves>stem>seeds. *M .azedarach* had more allelopathic effect than *T.orientalis* L. Results showed that both plants contain allelochemicals which show inhibitory effect against *P.ammericanum*.

5. Conclusion

From the present study it is concluded that *Meliaazedarach* L. and *Thujaorientalis* L. have allelochemicals which are secondary metabolites. These chemicals inhibit the seed germination, fresh and dry weight and seedling growth of *Pennisetumamericanum* L by affecting their respiration, cell division and different metabolic activities. *M.azedarach* L. has strong allelopathic potential. Plants show inhibitory effect due to water soluble allelochemicals present in their aqueous extracts. It can be used as herbicides for weed management

References

- [1] Alagesaboopathi, C. 2010. Allelopathic effect of *Centellaasiatica* aqueous extracts on Pearl millet (*Pennisetumtyphoides* L.) and Cowpea (*Vignaunguiculata* WALP.). Pak. J. Weed Sci. Res. 16(1): 67-71.
- [2] Atallah F. Mekhlif.2009. Effect of *Meliaazedarach* L. and *Ailanthus altissima*Swingle Extracts on the Larva Alimentary Tract and Growth of Black Cutworm, *Agrotisipsilon*Hufn. (Lepidoptera: Noctuidae). J. Raf. Sci., Vol. 20, No.2, pp 8- 18.
- [3] Fritz, D. A.P., J.S.Bernardi., B.M. Haas.,S.A.D.L.Ascoli., Bordignon, G.V. Bordignon., Poser.2007. Germination and growth in-

- hibitory effects of *Hypericum* and *H. polyanthemum* extracts on *Lactuca sativa* L, *Revista Brasileira de Farmacognosia. Brazilian Journal of Pharmacognosy*, 17 (1): 44-48.
- [4] Hussain, F., B., Ahmed and I. Ilahi. 2010. Allelopathic effects of *Cenchrus ciliaris* L. and *Bothriochloa pertusa* (L.) A. Camus. *Pak. J. Bot.*, 42(5): 3587-3604.
- [5] Khan, M., Hussain, F., Musharaf, S. (2011). Allelopathic potential of *Rhazya stricta* Decne on germination of *Pennisetum typhoides*. *International Journal of Biosciences (IJB)*, 1 (4): 80-85.
- [6] Isfahan, M.N., M, Shariat. 2007. The effect of some allelochemicals on seed germination of *Coronilla varia* L. seeds. *American-Eurasian J. Agric. & Environ. Sci*, 2 (5): 534-538.
- [7] Lungu, L., C.L, Popa, J, Morris and M, Savoiu. 2011. Evaluation of phytotoxic activity of *Melissaezardach* L. extracts on *Lactuca sativa* L. *Romanian Biotechnological Letters*, 16:(2)
- [8] Oyun, M.B. 2006. Allelopathic potentialities of *Gliricidia sepium* and *Acacia auriculiformis* on the germination and seedling vigour of Maize (*Zea mays* L.) *American journal of agricultural and biological science* 1 (3): 44-47. <http://dx.doi.org/10.3844/ajabssp.2006.44.47>.
- [9] Raja Nagappan. 2012. Impact of *Melissaezardach* Linn. (Meliaceae) Dry Fruit Extract, Farmyard Manure and Nitrogenous Fertilizer Application against Cabbage Aphid *Brevicoryne brassicae* Linn. (Homoptera: Aphididae) in Home Garden, *Asian Journal of Agricultural Sciences* 4(3): 193-197.
- [10] Siddiqui, S, S Bhardwaj, S.S. Khan and M.K. Meghvanshi. 2009. Allelopathic effect of different concentration of water extract of *Prosopis juliflora* leaf on seed germination and radical length of Wheat (*Triticum aestivum* Var-Lok-1). *American-Eurasian Journal of Scientific research*, 4(2): 81-84.
- [11] Singh, H.P, R.K, Kohli, D.R., Batish, P.S. Kaushal. 1999. Allelopathy of Gymnospermous trees. *J. For. Res.*, (4): 245-254. <http://dx.doi.org/10.1007/BF02762256>.
- [12] Smith, M.W., M.E., Wolf, B.S., Cheary and M.W. Carroll, Allelopathy of Bermudagrass, Tall Fescue, Redroot pigweed, and Cutleaf evening primrose on Pecan. *Hort Science*, 36(6):1047-1048.
- [13] Samreen, u., Hussain, F., Sher, Z. (2009). Allelopathic potential of *Calotropis procera* (AIT.) AIT. *Pak. J. Pl. Sci.*, 15 (1): 7-14.
- [14] Tadele, D and D, Teketay. 2010. Effects of extracts from leaves of *Eucalyptus globulus* Labill on seed germination and earl growth of *Olea europaea* L. subsp. *cuspidata*. CONFUSE.
- [15] Terzi, I. 2008. Allelopathic effects of Juglone and decomposed walnut leaf juice on muskmelon and cucumber seed germination and seedling growth. *African Journal of Biotechnology* Vol. 7 (12), pp. 1870-1874.
- [16] Terzi, I and I. Kocaçaliskan. 2009. Alleviation of juglone stress by plant growth regulators in germination of cress seeds. *Scientific Research and Essay* Vol. 4 (5) pp. 436-439.
- [17] Wirat Phuwawat, Watcharin Wichittrakarn, Chamroon Laosinwattana and Montinee Teerarak. 2012. Inhibitory Effects Of *Melissaezardach* L. Leaf Extracts On Seed Germination And Seedling Growth Of Two Weed Species, *Pak. J. Weed Sci. Res.*, 18: 485-492.