



# Planar and Non Planar Construction of $\gamma$ - Uniquely Colorable Graph

A. Elakkiya<sup>1</sup>, M. Yamuna<sup>2\*</sup>

<sup>1</sup>Vellore Institute of Technology, Vellore

<sup>2</sup> Vellore Institute of Technology, Vellore

\*Corresponding author E-mail: myamuna@vit.ac.in

## Abstract

A uniquely colorable graph  $G$  whose chromatic partition contains atleast one  $\gamma$  - set is termed as a  $\gamma$  - uniquely colorable graph. In this paper, we provide necessary and sufficient condition for  $\bar{G}$  and  $G^*$  to be  $\gamma$  - uniquely colorable whenever  $G$   $\gamma$ - uniquely colorable and also provide constructive characterization to show that whenever  $G$  is  $\gamma$ - uniquely colorable such that  $|P| \geq 2$ ,  $G$  can be both planar and non planar.

**Keywords:** Complement; Dual; Non Planar; Planar; Uniquely colorable graphs.

## 1. Introduction

In [1] Bing Zhou investigated the dominating  $\chi$ -color number,  $d_\chi(G)$ , of a graph  $G$ . In [2],[3], M. Yamuna et al introduced  $\gamma$  - uniquely colorable graphs and also provided the constructive characterization of  $\gamma$  - uniquely colorable trees and characterized planarity of complement of  $\gamma$  - uniquely colorable graphs. In [4],[5],M. Yamuna et al introduced Non domination subdivision stable graphs (NDSS) and characterized planarity of complement of NDSS graphs

## 2. Terminology

We consider simple graphs  $G$  with  $n$  vertices and  $m$  edges.  $K_n$  is a complete graph with  $n$  vertices.  $K_5$  and  $K_{3,3}$  are called Kuratowski's graph. Results related to graph theory we refer to [6].

Chromatic partition of a graph  $G$  is partition the vertices into smallest possible number of disjoint, independent sets. A graph  $G = (V, E)$  is said to be uniquely colorable if has a unique chromatic partition.

$D$  is dominating set if every vertex of  $V - D$  is adjacent to some vertex of  $D$ . Minimum cardinality of  $D$ , is said to be a minimum dominating set (MDS). The cardinality of any MDS for  $G$  is said to be domination number of  $G$ , represented by  $\gamma(G)$ . Results related to domination we refer to [7].

## 3. Result and Discussion

A uniquely colorable graph  $G$  whose chromatic partition contains atleast one  $\gamma$ - set is termed as a  $\gamma$ - uniquely colorable graph.

In Fig. 1  $G_1$  and  $G_2$  are  $\gamma$ - uniquely colorable graphs.  $G_1^*$  is  $\gamma$ - uniquely colorable while  $\bar{G}_2$  is not  $\gamma$ - uniquely colorable graph. So when  $G$  is  $\gamma$ - uniquely colorable,  $\bar{G}$  need not be  $\gamma$ - uniquely colorable. In Fig. 2  $G_1$  and  $G_2$  are  $\gamma$ - uniquely colorable graphs.  $G_1^*$  is  $\gamma$ - uniquely colorable while  $G_2^*$  is not  $\gamma$ - uniquely colorable

graph. So when  $G$  is  $\gamma$  - uniquely colorable,  $G^*$  need not be  $\gamma$ - uniquely colorable. In this paper, we determine the condition for  $\bar{G}$  and  $G^*$  to be  $\gamma$ - uniquely colorable whenever  $G$  is  $\gamma$ - uniquely colorable. We also provide the constructive characterization to show that whenever  $G$  is  $\gamma$  uniquely colorable such that  $|P| \geq 2$ ,  $G$  can be both planar and non planar.




Fig.1




Fig.2

**Theorem 1.** Let  $G$  be a  $\gamma$ -uniquely colorable graph.  $\bar{G}$  is also  $\gamma$ -uniquely colourable if and only if  $\exists$  a unique smallest possible partition  $P = \{ V_1, V_2, \dots, V_k \}$  of  $V(G)$   $\ni$

1. every  $V_i, i = 1$  to  $k$  is a clique
2. there exist one  $V_i \ni$  every vertex in  $V - \{ V_i \}$  is not adjacent to atleast one vertex in  $V_i$
3.  $|V_j| \geq |V_i|$ , for every  $i \neq j$
4.  $V_i$  is the smallest set in  $G$  satisfying 2

**Proof.** Assume that  $\bar{G}$  is a  $\gamma$ -uniquely colourable graph, implies there exist a partition  $P_1 = \{ V_1, V_2, \dots, V_k \}$  for  $\bar{G}$  such that

- $P_1$  is unique and smallest possible set.
- every  $V_i, i = 1$  to  $k$ , is independent in  $\bar{G}$  implies every  $V_i$  is a clique in  $G$ .
- there exist one  $V_i$  such that  $V_i$  is a  $\gamma$ -set for  $\bar{G}$ . Also  $|V_i| \geq |V_j|$ , for every  $i \neq j$  implies there exist one  $V_i$  in  $G$  such that every vertex in  $V - \{ V_i \}$  not adjacent to atleast one vertex in every  $V_i$ .

implies  $P_1 = \{ V_1, V_2, \dots, V_k \}$  is a  $\gamma$ -chromatic partition for  $V(G)$ .  $P_1$  is not unique implies  $\exists$  one  $P_2 = \{ W_1, W_2, \dots, W_k \}$  in  $G$  such that  $\{ W_1, W_2, \dots, W_k \}$  is a clique, implies  $P_2$  is also a  $\gamma$ -chromatic partition in  $\bar{G}$  such that every  $W_i$  is independent and  $|P_1| = |P_2|$ , a contradiction to our assumption that  $P_1$  is unique.  $P_1$  is not smallest, implies one  $P_3 = \{ V_1, V_2, \dots, V_k \}, q < k$  such that  $P_3$  is a  $\gamma$ -chromatic partition in  $G \ni U_i, i = 1$  to  $q$  is clique implies  $P_3$  is a  $\gamma$ -chromatic partition in  $\bar{G} \ni$  every  $U_i$  is independent and  $|P_3| < |P_1|$ , a contradiction.  $P_1$  is a  $\gamma$ -uniquely colorable partition for  $\bar{G}$  implies there exist one  $V_i$  such that  $V_i$  is a  $\gamma$ -set for  $\bar{G}$ , implies every vertex in  $V - \{ V_i \}$  is adjacent to atleast one vertex in  $V_i$ , implies  $P_1$  is a  $\gamma$ -chromatic partition in  $G \ni$  every vertex in  $V - \{ V_i \}$  is not adjacent to atleast one vertex in  $V_i$ . Also, we know that  $|V_i| \leq |V_j|$  for every  $i \neq j$  in  $\bar{G}$ , implies it is true in  $G$  also. If  $V_i$  is not the smallest set  $\ni$  every vertex in  $V - \{ V_i \}$  is  $\perp$  to atleast one vertex in  $V_i$  in  $\bar{G}$ , implies there exist one  $W$  contained in  $V(\bar{G})$  such that  $|W| < |V_i|$  and every vertex in  $W - V(\bar{G})$  is  $\perp$  to atleast one vertex in  $W$ , a contradiction  $\Rightarrow V_i$  is the smallest set satisfying the property. Hence  $P_1$  is a  $\gamma$ -chromatic partition in  $G \ni$  the conditions of the theorem are satisfied.

Conversely assume that the conditions of the theorem are satisfied.  $P$  is a partition such that it is unique and smallest such that every  $V_i$  is a clique, implies  $P_1$  is a partition in  $\bar{G}$  such that every  $V_i$  is independent. If  $P$  is not a smallest possible partition in  $\bar{G}$  then there exist one partition  $P_4 = \{ R_1, R_2, \dots, R_q \}, q < k$  in  $\bar{G}$  such that each  $R_i$  is independent, implies  $P_4$  is a partition in  $G$  such that every  $R_i$  is a clique such that  $|P_4| < |P|$ , a contradiction.  $P$  is not unique in  $\bar{G}$ , implies there exist a partition  $P_5 = \{ S_1, S_2, \dots, S_k \}$  such that each  $S_i$  is independent in  $\bar{G}$ , implies  $P, P_5$  are two possible partition with the same cardinality in  $G$ , a contradiction.  $P$  is a partition  $\ni$  there exist one  $V_i$ , every vertex in  $V - \{ V_i \}$  is not  $\perp$  to atleast one  $V_i, |V_j| \geq |V_i|$  for any  $i \neq j$  implies  $P$  is a partition in  $\bar{G} \ni$  every vertex in  $V - \{ V_i \}$  is adjacent to atleast one  $V_i, |V_i| \leq |V_j|, i \neq j$ , implies  $V_i$  is a dominating set for  $\bar{G}$ . Since  $V_i$  is the smallest set satisfying this property, implies  $V_i$  is the  $\gamma$ -set for  $\bar{G}$

Let  $P = \{ R_1, R_2, \dots, R_q \}$ , be the set of regions of  $G$ . Let  $T = \{ r_1, r_2, \dots, r_q \}$ , be the set of vertices in the regions  $R_1, R_2, \dots, R_q$  respectively, that is  $r_1$  is the vertex in the region  $R_1, r_2$  is the vertex in the region  $R_2, \dots, r_q$  is the vertex in the region  $R_q$  respectively.

We observe that

- There is a 1-1 mapping between  $S$  and  $T$ , i.e.  $\forall R_i \in S \exists r_i \in T, i = 1, \dots, q$ .
- $\forall X \subseteq S \exists$  a corresponding set in  $T$  (say  $X^*$ ), i.e. if  $X \subseteq S = \{ R_i, R_p, R_j \}$ , then  $X \subseteq T = \{ r_i, r_p, r_j \}$ .
- If  $a$  is any edge in  $G$  there is a corresponding edge in  $G^*$  (say  $a^*$ ).
- Let  $D \subseteq S \ni$  every region in  $S - D$  is  $\perp$  to atleast one region in  $D \Rightarrow \exists D^* \subseteq T \ni$  any vertex in  $T - D^*$  is  $\perp$  to atleast vertex in  $D^*$ .
- $D$  is a smallest cardinality satisfying this property  $\Rightarrow D^*$  is a  $\gamma$ -set for  $G^*$ .

**Theorem 2.** Let  $G$  be a  $\gamma$ -uniquely colourable graph.  $G^*$  is also  $\gamma$ -uniquely colourable graph if and only if there exist a unique smallest partition  $P = \{ R_1, R_2, \dots, R_k \}$  of  $R(G)$  such that

1. every  $R_i, i = 1, 2, \dots, k$  is independent.
2. there exist one  $R_i$  such that every region in  $R - \{ R_i \}$  is adjacent to atleast one region in  $R_i$ .
3.  $|R_j| \geq |R_i|$ .

**Proof.** Assume that  $G^*$  is  $\gamma$ -uniquely colourable graph. If  $G^*$  is  $\gamma$ -uniquely colourable graph, then there exist a partition  $P = \{ V_1, V_2, \dots, V_k \}$  such that  $P$  is a  $\gamma$ -chromatic partition,  $\Rightarrow$

1. every  $V_i$  is independent.
2.  $V_i$  is a  $\gamma$ -set for  $G^*$ .

1 implies, there exist a set of regions  $R_1, R_2, \dots, R_k$  in  $G$  such that every  $R_i$  is independent.

2 implies, there exist  $R_i \ni$  every region in  $R - \{ R_i \}$  is adjacent to atleast one region in  $R_i$  and  $R_i$  is the smallest set satisfying this property implies the conditions of the theorem are satisfied.

Conversely, assume that the conditions of the theorem are satisfied.  $P = \{ V_1, V_2, \dots, V_k \}$  is a partition of  $R(G)$ , implies there exist a partition  $P_1 = \{ V_1, V_2, \dots, V_k \}$  of  $V(G^*)$ .

- 1 implies, every  $V_i, i = 1, 2, \dots, k$  is independent.
- 2 implies, there exist one  $V_i \ni$  every vertex in  $V - \{ V_i \}$  is  $\perp$  to atleast one region in  $V_i$ .
- 3 implies  $|V_j| \geq |V_i|$  for all  $i \neq j$

Since  $P$  is a unique partition there exist no other partition of  $V(G^*)$  that satisfies all these conditions implies,  $P_1$  is a  $\gamma$ -chromatic partition for  $G^*$ .

**Planar and Non planar Construction**

In this section, we provide constructive characterization to show that whenever  $G$  is  $\gamma$ -uniquely colorable such that  $|P| \geq 2, G$  can be both planar and nonplanar.

Planar Construction when  $|P| = 2$ .

Let  $\gamma(G) = k_1$ . Let  $P = \{ V_1, V_2 \}$ , where  $V_1 = \{ a_1, a_2, \dots, a_{k_1} \}$   $V_2 = \{ b_1, b_2, \dots, b_{k_2} \}, k_2 \geq k_1, k_1 \geq 3, k_2 \geq 4$ .

Construct a graph  $G_1$  as follows

1.  $V(G_1) = V(G)$
2. Consider  $k_1$  vertices in  $V_1$  and  $V_2$  say  $\{ a_1, a_2, \dots, a_{k_1} \}$  and  $\{ b_1, b_2, \dots, b_{k_2} \}$ .

Construct a comb graph with  $2k_1$  vertices. Label the vertices of this comb as seen in Fig. 3




Fig.3

Include the remaining  $k_2 - k_1$  vertices of  $V_2$  as pendant vertices with  $a_{k_1}$  as the support vertex. The general structure of graph  $G_1$  is as seen in the Fig.4.




Fig.4

Since we have atleast  $k_1$  pendant vertices,  $\gamma(G_1) \geq k_1$ .  $\{a_1, a_2, \dots, a_{k_1}\}$  is a dominating set for  $G_1$ , implies  $\gamma(G_1) = k_1$ . Since  $\langle a_1, a_2, \dots, a_{k_1}, b_1, b_2, \dots, b_{k_2} \rangle$  is acomb, the only possible maximal independent sets are  $\{a_1, a_2, \dots, a_{k_1}\}$  and  $\{b_1, b_2, \dots, b_{k_2}\}$ .  $P = \{V_1, V_2\}$  is a partition for  $G_1$  such that

1.  $V_1$  is  $\gamma$ - set for  $G_1$
2.  $P$  is the only possible partition for  $G_1$ ,  $\Rightarrow G_1$  is a  $\gamma$ - uniquely colorable graph.

Non Planar Construction when  $|P| = 2$ .

Let  $\gamma(G) \geq k_1$ ,  $k_1 \geq 6$ ,  $P = \{V_1, V_2\}$   $V_1 = \{a_1, a_2, \dots, a_{k_1}\}$ ;  $V_2 = \{b_1, b_2, \dots, b_{k_2}\}$ ,  $k_2 \geq 6$

Construct a graph  $G_1$  as follows

1.  $V(G_1) = V(G)$
2. Consider  $k_1$  vertices in  $V_1$  and  $k_2$  vertices in  $V_2$  say  $\{a_1, a_2, a_3, a_4, a_5, a_6\}$ ,  $\{b_1, b_2, b_3, \dots, b_{k_2}\}$ . Let  $\langle a_1, a_2, a_3, b_1, b_2, b_3 \rangle$  is  $K_{3,3}$ . Include the remaining  $a_i, b_i, i = 1, 2, 3$ . Include the remaining  $b_i, i = 6, 7, \dots, k_2$  as arbitrary pendant vertices adjacent to any  $a_i, i = 1, 2, 3$ .
3. Graph  $G_1$  is as seen in Fig.5.




Fig.5.

Since  $G_1$  has atleast  $k_1$  pendant vertices  $\{a_4, a_5, \dots, a_{k_1}, b_4, b_5, \dots, b_{k_1}\}$ ,  $\gamma(G_1) \geq k_1$ ,  $\{V_1\}$  dominates  $G_1$ . Also  $|V_1| = k_1$ , implies that  $V_1$  is  $\gamma$ - set for  $G_1$ , since  $G_1$  is a bipartite graph  $P = \{V_1, V_2\}$  is the only chromatic partition for  $G_1$  such that  $V_1$  is a  $\gamma$ - set for  $G_1$ , implies  $G_1$  is  $\gamma$ - uniquely colorable and non planar.

$\gamma(G) = 3$ ,  $P = \{V_1, V_2\}$ ,  $V_1 = \{a_1, a_2, a_3\}$ ,  $V_2 = \{b_1, b_2, \dots, b_{k_1}\}$ ,  $k_1 \geq 6$ ,  
 $\gamma(G) = 4$ ,  $P = \{V_1, V_2\}$ ,  $V_1 = \{a_1, a_2, a_3, a_4\}$ ,  $V_2 = \{b_1, b_2, \dots, b_{k_1}\}$ ,  $k_1 \geq 6$ ,  
 $\gamma(G) = 5$ ,  $P = \{V_1, V_2\}$ ,  $V_1 = \{a_1, a_2, a_3, a_4, a_5\}$ ,  $V_2 = \{b_1, b_2, \dots, b_{k_1}\}$ ,  $k_1 \geq 6$ , are analogous to the above discussion.

$|P| = 3 = P = \{V_1, V_2, V_3\}$ ,  $|V_1| = k_1$ ,  $|V_2| = k_2$ ,  $|V_3| = k_3$ ,  $k_2, k_3 \geq k_1$ . Planar Construction when  $|P| = 3$ .

$|P| = 3 = P = \{V_1, V_2, V_3\}$ ,  $|V_1| = k_1$ ,  $|V_2| = k_2$ ,  $|V_3| = k_3$ ,  $k_2, k_3 \geq k_1$ . Consider a wheel graph with  $k$  vertices where  $k = k_1 + 2k_i$ , where  $k_i = \min(k_2, k_3)$ . Label the vertices of the wheel in the following fashion as seen in Fig.6.




Fig.6

If  $k_2 \neq k_3$ , then we include the remaining vertices as follows.

Let  $k_2 > k_3$ . Let  $k_2 = k_3 + m$ . Label the additional vertices as  $\{b_{k_3+1}, b_{k_3+2}, \dots, b_{k_2}\}$ . Include these vertices as seen in Fig.7.




Fig.7

Since  $\langle b_1, a_j, c_i \rangle, i = 1$  to  $k_1 - 1, j = 2$  to  $a_{k_1}$  is  $P_3$  either  $a_j$  or  $b_1$  or  $c_i$  should be included in every possible  $\gamma$ - set for  $G$ .  $\{a_1, a_2, \dots, a_{k_1}\}$  is a  $\gamma$ - set for  $G$ . Also  $\{V_1, V_2, V_3\}$  is the only possible chromatic partition for  $G$  implies  $\gamma$ - uniquely colorable graph  $G$  is planar.

### 4. Conclusion

In this paper, we provide necessary and sufficient condition for  $\bar{G}$  and  $G^*$  to be  $\gamma$  uniquely colorable and also provide constructive characterization to show that whenever  $G$  is  $\gamma$ - uniquely colorable such that  $|P| \geq 2$ ,  $G$  can be both planar and non planar.

### References

- [1] Bing Zhou, "On the maximum number of dominating classes in graph coloring", *Open Journal of Discrete Mathematics*, Vol 6, (2016), pp.70 – 73.
- [2] M. Yamuna, A. Elakkiya, "  $\gamma$  - Uniquely colorable graphs", *IOP Conf. Series: Materials Science and Engineering*, Vol.263, (2017).
- [3] M. Yamuna, A. Elakkiya, " Planar graph characterization of  $\gamma$ - Uniquely colorable graphs", *IOP Conf. Series: Materials Science and Engineering*, Vol.263, (2017).
- [4] Yamuna, M., Elakkiya, A., "Non domination subdivision stable graphs", *IOP Conf. Series: Materials Science and Engineering*. Vol 263, (2017).
- [5] Yamuna, M., Elakkiya, A., "Planar graph characterization of NDSS graphs", *IOP Conf. Series: Materials Science and Engineering*, Vol 263, (2017).
- [6] Harary, F., *Graph Theory*, Addison Wesley, Narosa Publishing House, (2001).
- [7] Haynes, T.W., Hedetniemi, S. T & Slater, P. J. *Fundamentals of domination in graphs*, New York, Marcel Dekker, (1998).